

30009557 RECEIVED
 RECEIVED PCT/PTO 08 NOV 2001

FORM PTO-1390 OFFICE		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK	ATTORNEY'S DOCKET NUMBER PF-0693 USN
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371			U.S. APPLICATION NO. (If known, see 37 CFR 1.5) TO BE ASSIGNED 10/009557
INTERNATIONAL APPLICATION NO PCT/US00/12811	INTERNATIONAL FILING DATE 10 May 2000	PRIORITY DATE CLAIMED 11 May 1999	
TITLE OF INVENTION EXTRACELLULAR MATRIX AND ADHESION-ASSOCIATED PROTEINS			
APPLICANT(S) FOR DO/EO/US BANDMAN, Olga; HILLMAN, Jennifer L.; TANG, Y. Tom; LAL, Preeti; YUE, Henry; BAUGHN, Mariah R.; LU, Dyung Aina M.; AZIMZAI, Yalda			
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: 1. <input checked="" type="checkbox"/> This is the FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input type="checkbox"/> This is an express request to promptly begin national examination procedures (35 U.S.C. 371 (f)). 4. <input type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (PCT Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau) b. <input type="checkbox"/> has been communicated by the International Bureau. c. <input checked="" type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).			
Items 11 to 16 below concern document(s) or information included: 11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input checked="" type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.27 and 3.31 is included. 13. <input checked="" type="checkbox"/> A FIRST preliminary amendment, as follows: Cancel in this application original claims #16, 19, and 22 before calculating the filing fee, without prejudice or disclaimer. Applicants submit that these claims were included in the application as filed in the interest of providing notice to the public of certain specific subject matter intended to be claimed, and are being canceled at this time in the interest of reducing filing costs. Applicants expressly state that these claims are not being canceled for reasons related to patentability, and are in fact fully supported by the specification as filed. Applicants expressly reserve the right to reinstate these claims or to add other claims during prosecution of this application or a continuation or divisional application. Applicants expressly do not disclaim the subject matter of any invention disclosed herein which is not set forth in the instantly filed claims. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 14. <input type="checkbox"/> A substitute specification. 15. <input type="checkbox"/> A change of power of attorney and/or address letter. 16. <input checked="" type="checkbox"/> Other items or information: 1) Transmittal Letter (2 pp, in duplicate) 2) Return Postcard 3) Express Mail Label No.: EL 856 148830 US 4) Request to Transfer			

JC13 Rec'd PCT/PTC 08 NOV 2001

U.S. APPLICATION NO. (if known, see 37 CFR 1.5) TO BE ASSIGNED 10/009557	INTERNATIONAL APPLICATION NO PCT/US00/12811	ATTORNEY'S DOCKET NUMBER PF-0693 USN
17. <input checked="" type="checkbox"/> The following fees are submitted BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO..... \$1000.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO..\$860.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$710.00 <input checked="" type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4).....\$690.00 <input type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) .. \$100.00		
ENTER APPROPRIATE BASIC FEE AMOUNT =		\$710.00
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e))		\$
CLAIMS	NUMBER FILED	NUMBER EXTRA
Total Claims	20 =	0
Independent Claims	2 =	0
MULTIPLE DEPENDENT CLAIM(S) (if applicable)		+ \$270.00
TOTAL OF ABOVE CALCULATIONS =		\$
<input type="checkbox"/> Applicant claims small entity status See 37 CFR 1.27. The fees indicated above are reduced by 1/2.		\$
SUBTOTAL		\$710.00
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f))		\$
TOTAL NATIONAL FEE =		\$710.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)) The assignment must be accompanied by the appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property		\$
TOTAL FEES ENCLOSED =		\$710.00
		Amount to be Refunded:
		Charged
a. <input type="checkbox"/> A check in the amount of \$_____ to cover the above fees is enclosed. b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. <u>09-0108</u> in the amount of \$710.00 _____ to cover the above fees. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>09-0108</u> . A duplicate copy of this sheet is enclosed		
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.		
SEND ALL CORRESPONDENCE TO INCYTE GENOMICS, INC 3160 Porter Drive Palo Alto, CA 94304		
 SIGNATURE		
NAME: Diana Hamlet-Cox		
REGISTRATION NUMBER 33,302		
DATE: <u>8</u> November 2001		

Certificate of Mailing

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Commissioner for Patents, Washington, D.C. 20231 on July 16 2002.

By: Christopher R. Leach Printed: Christopher R. Leach

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: **Bandman et al.**

Title: **EXTRACELLULAR MATRIX AND ADHESION-ASSOCIATED PROTEINS**

Serial No.: **10/009,557**

Filing Date: **To Be Assigned**

Examiner: **To Be Assigned**

Group Art Unit: **To Be Assigned**

Box PCT
Commissioner for Patents
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Sir:

Prior to examination, please amend the above-referenced application as follows:

IN THE CLAIMS

If claims 24-95 are in fact still pending in this application, cancel claims 24-95, prior to examination of the claims, without prejudice or disclaimer.

Attached hereto is a marked-up version of the changes made to the claims by the current amendment. The attached page is captioned "Version with markings to show changes made."

Docket No.: PF-0693 USN

REMARKS

Applicants would like to verify that the above claims 24-95 are in fact canceled prior to examination of the instant application, but **after** their inclusion in the specification as filed, so that they will be published.


Applicants respectfully submit that failure to clearly cancel these claims on the Transmittal Sheet for the National Stage application was an oversight, and that the Article 34 Amendment was overlooked when preparing the patent application Transmittal Sheet.

Applicants note that it was their intention to have the above-referenced claims published with the United States application in the interest of providing notice to the public of certain specific subject matter intended to be claimed, and were to be canceled prior to examination in the interest of reducing filing costs at the present time. Applicants expressly state that these claims were not being canceled for reasons related to patentability, and are in fact fully supported by the specification as filed. Applicants expressly reserve the right to reinstate these claims or to add other claims during prosecution of this application or a continuation or divisional application. Applicants expressly do not disclaim the subject matter of any invention disclosed herein which is not set forth in the instantly filed claims.

Applicants believe that no fee is due with this communication. However, if the USPTO determines that a fee is due, the Commissioner is hereby authorized to charge Deposit Account No. **09-0108**, as set forth in the enclosed fee transmittal letter. Applicants further request that any refund that might be due to Applicants be credited to Deposit Account No. **09-0108** as well.

Respectfully submitted,

INCYTE GENOMICS, INC.

Date: 10 July 2002

Diana Hamlet-Cox

Reg. No. 33,302

Direct Dial Telephone: (650) 845-4639

3160 Porter Drive
Palo Alto, California 94304
Phone: (650) 855-0555
Fax: (650) 849-8886

Jul 17, 2002 4:26PM INCYTE LEGAL DEPT

ALCANTARA No. 360921-0004

Docket No.: PF-0693 USN

VERSION WITH MARKINGS TO SHOW CHANGES MADE

IN THE CLAIMS

Claim(s) 24-95 have been canceled without prejudice or disclaimer.

EXTRACELLULAR MATRIX AND ADHESION-ASSOCIATED PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of extracellular matrix and
5 adhesion-associated proteins and to the use of these sequences in the diagnosis, treatment, and
prevention of cell proliferative, immune, reproductive, neuronal, and genetic disorders.

BACKGROUND OF THE INVENTION

Extracellular Matrix Proteins

10 The extracellular matrix (ECM) is a complex network of glycoproteins, polysaccharides,
proteoglycans, and other macromolecules that are secreted from the cell into the extracellular space.
The ECM remains in close association with the cell surface and provides a supportive meshwork that
profoundly influences cell shape, motility, strength, flexibility, and adhesion. In fact, adhesion of a cell
to its surrounding matrix is required for cell survival except in the case of metastatic tumor cells, which
15 have overcome the need for cell-ECM anchorage. This phenomenon suggests that the ECM plays a
critical role in the molecular mechanisms of growth control and metastasis. (Reviewed in Ruoslahti, E.
(1996) Sci. Am. 275:72-77.) Furthermore, the ECM determines the structure and physical properties
of connective tissue and is particularly important for morphogenesis and other processes associated
with embryonic development and pattern formation.

20

Collagens

The collagens comprise a family of ECM proteins that provide structure to bone, teeth, skin,
ligaments, tendons, cartilage, blood vessels, and basement membranes. Multiple collagen proteins have
been identified. Three collagen molecules fold together in a triple helix stabilized by interchain disulfide
25 bonds. Bundles of these triple helices then associate to form fibrils. Collagen primary structure
consists of hundreds of (Gly-X-Y) repeats where about a third of the X and Y residues are Pro.
Glycines are crucial to helix formation as the bulkier amino acid side chains cannot fold into the triple
helical conformation. Because of these strict sequence requirements, mutations in collagen genes have
severe consequences. Osteogenesis imperfecta patients have brittle bones that fracture easily; in severe
30 cases patients die in utero or at birth. Ehler-Danlos syndrome patients have hyperelastic skin,
hypermobile joints, and susceptibility to aortic and intestinal rupture. Chondrodysplasia patients have
short stature and ocular disorders. Alport syndrome patients have hematuria, sensorineural deafness,
and eye lens deformation. (See Isselbacher, K.J., et al. (1994) Harrison's Principles of Internal
Medicine, McGraw-Hill, Inc., New York, NY, pp. 2105-2117; and Creighton, T.E. (1984) Proteins,

WO 00/68380

PCT/US00/12811

cross by disulfide bonds. Laminin is especially important for angiogenesis and, in particular, for guiding the formation of capillaries. (Reviewed in Alberts, supra, pp. 990-991.)

Proteoglycans

- 5 There are many other types of proteinaceous ECM components, most of which can be classified as proteoglycans. Proteoglycans are composed of unbranched polysaccharide chains (glycosaminoglycans) attached to protein cores. Common proteoglycans include aggrecan, betaglycan, decorin, perlecan, serglycin, and syndecan-1. Some of these molecules not only provide mechanical support, but also bind to extracellular signaling molecules, such as fibroblast growth factor and
- 10 transforming growth factor β , suggesting a role for proteoglycans in cell-cell communication. (Reviewed in Alberts, supra, pp. 973-978.) Likewise, the glycoproteins tenascin-C and tenascin-R are expressed in developing and lesioned neural tissue and provide stimulatory and anti-adhesive (inhibitory) properties, respectively, for axonal growth (Faissner, A. (1997) Cell Tissue Res. 290:331-341).
- 15 Dentin phosphoryn (DPP) is a major component of the dentin ECM. DPP is a proteoglycan that is synthesized and expressed by odontoblasts (Gu, K., et al. (1998) Eur. J. Oral Sci. 106:1043-1047). DPP is believed to nucleate or modulate the formation of hydroxyapatite crystals. The gene encoding DPP has been mapped to human chromosome 4. Chromosome 4 contains the gene loci for two dentin genetic diseases, dentinogenesis imperfecta type II and dentin dysplasia type II (Feng, J.Q.,
- 20 et al. (1998) J. Biol. Chem. 273:9457-9464).

Mucins

- Mucins are highly glycosylated glycoproteins that are the major structural component of the mucus gel. The physiological functions of mucins are cytoprotection, mechanical protection,
- 25 maintenance of viscosity in secretions, and cellular recognition. MUC6 is a human gastric mucin that is also found in gall bladder, pancreas, seminal vesicles, and female reproductive tract (Toribara, N.W., et al. (1997) J. Biol. Chem. 272:16398-16403). The MUC6 gene has been mapped to human chromosome 11 (Toribara, N.W., et al. (1993) J. Biol. Chem. 268:5879-5885). Hemomucin is a novel Drosophila surface mucin that may be involved in the induction of antibacterial effector molecules
- 30 (Theopold, U., et al. (1996) J. Biol. Chem. 271:12708-12715).

Link Protein

Link protein binds to both cartilage proteoglycan and hyaluronan in cartilage ECM. This binding stabilizes the aggregation of these cartilage ECM proteins and produces supramolecular

WO 00/68380

PCT/US00/12811

assemblies. Link protein has been detected in other connective tissues, where it may bind proteoglycans and hyaluronan. Link protein contains a signal peptide, an immunoglobulin repeat, and link repeats (Ayad, S., et al. (1994) The Extracellular Matrix Facts Book, Academic Press, Inc., San Diego, CA, pp. 120-121).

5

Adhesion-Associated Proteins

The surface of a cell is rich in transmembrane proteoglycans, glycoproteins, glycolipids, and receptors. These macromolecules mediate adhesion with other cells and with components of the ECM. The interaction of the cell with its surroundings profoundly influences cell shape, strength, flexibility, 10 motility, and adhesion. These dynamic properties are intimately associated with signal transduction pathways controlling cell proliferation and differentiation, tissue construction, and embryonic development.

Cadherins

15 Cadherins comprise a family of calcium-dependent glycoproteins that function in mediating cell-cell adhesion in virtually all solid tissues of multicellular organisms. These proteins share multiple repeats of a cadherin-specific motif, and the repeats form the folding units of the cadherin ECM. Cadherin molecules cooperate to form focal contacts, or adhesion plaques, between adjacent epithelial cells. The cadherin family includes the classical cadherins and protocadherins. Classical cadherins 20 include the E-cadherin, N-cadherin, and P-cadherin subfamilies. E-cadherin is present on many types of epithelial cells and is especially important for embryonic development. P-cadherin is present on cells of the placenta and epidermis. Recent studies report that protocadherins are involved in a variety of cell-cell interactions (Suzuki, S. T. (1996) J. Cell Sci. 109:2609-2611). The intracellular anchorage of cadherins is regulated by their dynamic association with catenins, a family of cytoplasmic signal 25 transduction proteins associated with the actin cytoskeleton. The anchorage of cadherins to the actin cytoskeleton appears to be regulated by protein tyrosine phosphorylation, and the cadherins are the target of phosphorylation-induced junctional disassembly (Aberle, H., et al. (1996) J. Cell. Biochem. 61:514-523).

30 Integrins

Integrins are ubiquitous transmembrane adhesion molecules that link the ECM to the internal cytoskeleton. Integrins are composed of two noncovalently associated transmembrane glycoprotein subunits called α and β . Integrins function as receptors that play a role in signal transduction. For example, binding of integrin to its extracellular ligand may stimulate changes in intracellular calcium

5

10

15

25

30

Selectins

Selectins, or LEC-CAMs, comprise a specialized lectin subfamily involved primarily in inflammation and leukocyte adhesion. (Reviewed in Lasky, *supra*.) Selectins, which mediate the recruitment of leukocytes from the circulation to sites of acute inflammation, are expressed on the surface of vascular endothelial cells in response to cytokine signaling. Selectins bind to specific ligands on the leukocyte cell membrane and enable the leukocyte to adhere to and migrate along the endothelial surface. Binding of selectin to its ligand leads to polarized rearrangement of the actin cytoskeleton and stimulates signal transduction within the leukocyte (Brenner, B., et al. (1997) *Biochem. Biophys. Res. Commun.* 231:802-807; Hidari, K.I., et al. (1997) *J. Biol. Chem.* 272:28750-28756). Members of the selectin family possess three characteristic motifs: a lectin or carbohydrate recognition domain; an epidermal growth factor (EGF)-like domain; and a variable number of short consensus repeats (scr or "sushi" repeats) which are also present in complement regulatory proteins. The selectins include lymphocyte adhesion molecule-1 (LAM-1 or L-selectin), endothelial leukocyte adhesion molecule-1 (ELAM-1 or E-selectin), and granule membrane protein-140 (GMP-140 or P-selectin) (Johnston, G.I., et al. (1989) *Cell* 56:1033-1044).

Attractin

Attractin is a 134 kilodalton glycoprotein found in the serum. It is a member of the CUB family of cell adhesion proteins and binds directly to leukocytes. Attractin has a CUB domain, an EGF domain, and C-type lectin protein domains. This serum protein mediates the interaction between T lymphocytes and monocytes and leads to the adherence and spreading of monocytes that become the foci for T cell clustering. (See, Duke-Cohan, J.S., et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:11336-11341.)

25 Proteins Containing Leucine Rich Repeats (LRRs)

LRRs are sequence motifs, approximately 22-28 amino acids in length, found in proteins with a large variety of functions and cellular locations. Proteins containing LRRs are all thought to be involved in protein-protein interactions. The crystal structure of LRRs has been studied and found to correspond to beta-alpha structural units. These structural units form a parallel beta sheet with one surface exposed to solvent. In this way an LRR-containing protein acquires a nonglobular shape (Kobe, B. and Deisenhofer, J. (1994) *Trends Biochem. Sci.* 19:415-421). There is evidence to suggest LRRs function in signal transduction and cellular adhesion as well as in protein-protein interactions (Gay, N.J., et al. (1991) *FEBS Lett.* 29:87-91). For example, LLR proteins such as connectin and chaoptin are important cell adhesion molecules in neuronal development in *Drosophila melanogaster*.

and mammalian homologs are found in mouse (Taguchi, et al. (1996) Brain Res. Mol. Brain Res. 1-2:31-40).

Proteins Containing Armadillo/ β -Catenin-like Repeats

- 5 Various proteins such as those encoded by the Drosophila armadillo gene and the human APC gene contain amino acid repeats that interact with β -catenins. The armadillo gene is required for pattern formation within the embryonic segments and imaginal discs and is highly conserved. It is 63% identical to a human protein, plakoglobin, which is involved in adhesive junctions joining epithelial and other cells (Peifer, M. and Wieschaus, E. (1990) Cell 63:1167-1176). APC gene mutations appear to
- 10 initiate inherited forms of human colorectal cancer and sporadic forms of colorectal and gastric cancer (Rubinfeld, B., et al. (1993) Science 262:1731-1734). The fact that the protein encoded by APC interacts with catenin suggests a link between tumor initiation and cell adhesion (Su, L.K., et al. (1993) Science 262:1734-1737).

15 Proteins Containing C-type Lectin Domains

- C-type lectin domains are found in a variety of proteins, including selectins and lecticans. Lecticans are a family of chondroitin sulfate proteoglycans that include aggrecan, versican, neurocan, and brevican. All C-type lectin proteins are involved in protein-protein interactions (Aspberg, A., et al. (1997) Proc. Natl. Acad. Sci. USA 94:10116-10121). A novel macrophage-restricted C-type lectin
- 20 protein has been cloned from mouse tissue. It is a type II transmembrane protein with one extracellular C-type lectin domain (Balch, S.G., et al. (1998) J. Biol. Chem. 273:18656-18664).

Bystin

- Bystin is a cytoplasmic protein that binds directly to trophinin, a cell adhesion molecule, and
- 25 tastin. The three molecules form a complex that is involved in cell adhesion. Bystin, tastin, and trophinin are strongly expressed in cells involved in the implantation of embryos, specifically in cells at human implantation sites and in intermediate trophoblasts at the invasion front of the placenta in early pregnancy. Bystin also binds to cytokeratins. During early embryogenesis cytokeratins 8 and 18 are expressed in the trophectoderm of blastocytes. It is possible that the molecular complex formed by
- 30 bystin, tastin, and trophinin interacts with the cytokeratins of trophectoderm cells at the time of implantation. A key component of embryo implantation is the unique cell adhesion to endometrial epithelium that occurs and the subsequent invasion of the maternal tissue by the trophoblast. Bystin may have an important role in the signal transduction that links cell adhesion to proliferation (Suzuki, N., et al. (1998) Proc. Natl. Acad. Sci. 95:5027-5032).

The invention further provides an isolated polynucleotide encoding a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25. In one alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:26-50.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25.

The invention further provides an isolated polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50, b) a naturally occurring

polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50, c) a polynucleotide sequence complementary to a), or d) a polynucleotide sequence complementary to b). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

5 Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50, c) a polynucleotide sequence complementary to a), or
10 d) a polynucleotide sequence complementary to b). The method comprises a) hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and b) detecting the presence or absence of said hybridization complex, and
15 optionally, if present, the amount thereof. In one alternative, the probe comprises at least 30 contiguous nucleotides. In another alternative, the probe comprises at least 60 contiguous nucleotides.

 The invention further provides a pharmaceutical composition comprising an effective amount of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an
20 amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, and a pharmaceutically acceptable excipient. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional EXMAD, comprising
25 administering to a patient in need of such treatment the pharmaceutical composition.

 The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically
30 active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an agonist compound identified by the method and a

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing
5 particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a
10 reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to
15 practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

20 DEFINITIONS

"EXMAD" refers to the amino acid sequences of substantially purified EXMAD obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of
25 EXMAD. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of EXMAD either by directly interacting with EXMAD or by acting on components of the biological pathway in which EXMAD participates.

An "allelic variant" is an alternative form of the gene encoding EXMAD. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in
30 polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding EXMAD include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as EXMAD or a polypeptide with at least one functional characteristic of EXMAD. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding EXMAD, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding EXMAD. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent EXMAD. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of EXMAD is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where “amino acid sequence” is recited to refer to an amino acid sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

“Amplification” relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term “antagonist” refers to a molecule which inhibits or attenuates the biological activity of EXMAD. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of EXMAD either by directly interacting with EXMAD or by acting on components of the biological pathway in which EXMAD participates.

The term “antibody” refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind EXMAD polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired.

Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

5 The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

10 The term "antisense" refers to any composition capable of base-pairing with the "sense" strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-
15 methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can
20 refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic EXMAD, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

25 The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid
30 strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or

side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "fragment" is a unique portion of EXMAD or the polynucleotide encoding EXMAD which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:26-50 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:26-50, for example, as distinct from any other sequence in the same genome. A fragment of SEQ ID NO:26-50 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:26-50 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:26-50 and the region of SEQ ID NO:26-50 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-25 is encoded by a fragment of SEQ ID NO:26-50. A fragment of SEQ ID NO:1-25 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-25. For example, a fragment of SEQ ID NO:1-25 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-25. The precise length of a fragment of SEQ ID NO:1-25 and the region of SEQ ID NO:1-25 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended

purpose for the fragment.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequence pairs.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis

15

Filter: on

20

25

30

18

of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) with blastp set at default parameters. Such default parameters may be, for example:

10 *Matrix: BLOSUM62*
 Open Gap: 11 and Extension Gap: 1 penalties
 Gap x drop-off: 50
 Expect: 10
 Word Size: 3
 15 *Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number. or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term "humanized antibody" refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of identity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding

WO 00/68380

PCT/US00/12811

between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive
 5 annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Generally, such wash temperatures are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic
 10 strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook et al., 1989. Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY: specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance,
 20 denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a
 25 similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g.,
 30 paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune

WO 00/68380

PCT/US00/12811

sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis et al., 1990, PCR Protocols. A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary

WO 00/68380

PCT/US00/12811

polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence.

- 5 This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence.
- 10 Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

- An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear
- 15 sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

- The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding EXMAD, or fragments thereof, or EXMAD itself, may comprise a bodily fluid; an
- 20 extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

- The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure
- 25 of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

- The term "substantially purified" refers to nucleic acid or amino acid sequences that are
- 30 removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

5 "Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, 10 heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to 15 animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in 20 vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, and plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention 25 into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) 30 set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides 35 due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may

Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

15 The invention is based on the discovery of new human extracellular matrix and adhesion-associated proteins (EXMAD), the polynucleotides encoding EXMAD, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative, immune, reproductive, neuronal, and genetic disorders.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding EXMAD. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each EXMAD were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. In some cases, GenBank sequence identifiers are also shown in column 5. The Incyte clones and GenBank cDNA sequences, where indicated, in column 5 were used to assemble the consensus nucleotide sequence of each EXMAD and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

35 The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions

associated with nucleotide sequences encoding EXMAD. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:26-50 and to distinguish between SEQ ID NO:26-50 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express EXMAD as a fraction of total tissues expressing EXMAD. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing EXMAD as a fraction of total tissues expressing EXMAD. Column 5 lists the vectors used to subclone each cDNA library.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding EXMAD were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

SEQ ID NO:42 maps to chromosome 8 within the interval from 64.60 to 90.20 centiMorgans.

SEQ ID NO:48 maps to chromosome 2 within the interval from 193.60 to 197.60 centiMorgans.

The invention also encompasses EXMAD variants. A preferred EXMAD variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the EXMAD amino acid sequence, and which contains at least one functional or structural characteristic of EXMAD.

The invention also encompasses polynucleotides which encode EXMAD. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:26-50, which encodes EXMAD. The polynucleotide sequences of SEQ ID NO:26-50, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding EXMAD. In particular, such a variant polynucleotide sequence will have at least about 80%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding EXMAD. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:26-50 which has at least about 80%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:26-50. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of EXMAD.

maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding EXMAD may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; and Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively, EXMAD itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of EXMAD, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins. Structures and Molecular Properties, WH Freeman, New York NY.)

In order to express a biologically active EXMAD, the nucleotide sequences encoding EXMAD or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding EXMAD. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding EXMAD. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding EXMAD and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) Results Probl. Cell Differ. 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding EXMAD and appropriate transcriptional and translational

WO 00/68380

PCT/US00/12811

control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding EXMAD. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding EXMAD. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding EXMAD can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding EXMAD into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of EXMAD are needed, e.g. for the production of antibodies, vectors which direct high level expression of EXMAD may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of EXMAD. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra: Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of EXMAD. Transcription of sequences encoding EXMAD may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J.

WO 00/68380

PCT/US00/12811

Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding EXMAD is inserted within a marker gene sequence, transformed cells containing sequences encoding EXMAD can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding EXMAD under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding EXMAD and that express EXMAD may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of EXMAD using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on EXMAD is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding EXMAD include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding EXMAD, or any fragments thereof, may be cloned into a vector

WO 00/68380

PCT/US00/12811

- salivary glands, skin, spleen, testis, thymus, thyroid, and uterus: an immune disorder, such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis.
- 5 autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome,
- 10 episodic lymphopenia with lymphocytotoxins, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis.
- 15 Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; a reproductive disorder, such as a disorder of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian
- 20 hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis: cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; a neuronal disorder, such as akathisia, Alzheimer's disease,
- 25 amnesia, amyotrophic lateral sclerosis, bipolar disorder, catatonia, cerebral neoplasms, dementia, depression, diabetic neuropathy, Down's syndrome, tardive dyskinesia, dystonias, epilepsy, Huntington's disease, peripheral neuropathy, multiple sclerosis, neurofibromatosis, Parkinson's disease, paranoid psychoses, postherpetic neuralgia, schizophrenia, and Tourette's disorder; and a genetic disorder, such as adrenoleukodystrophy, Alport's syndrome, choroideremia, Duchenne and
- 30 Becker muscular dystrophy, Down's syndrome, cystic fibrosis, chronic granulomatous disease, dentinogenesis imperfecta type II, dentin dysplasia type II, Gaucher's disease, Huntington's chorea, Marfan's syndrome, muscular dystrophy, myotonic dystrophy, pycnodysostosis, Refsum's syndrome, retinoblastoma, sickle cell anemia, thalassemia, Werner syndrome, von Willebrand's disease, Wilms' tumor, Zellweger syndrome, peroxisomal acyl-CoA oxidase deficiency, peroxisomal thiolase
- 35 deficiency, peroxisomal bifunctional protein deficiency, mitochondrial carnitine palmitoyl transferase

and carnitine deficiency, mitochondrial very-long-chain acyl-CoA dehydrogenase deficiency, mitochondrial medium-chain acyl-CoA dehydrogenase deficiency, mitochondrial short-chain acyl-CoA dehydrogenase deficiency, mitochondrial electron transport flavoprotein and electron transport flavoprotein:ubiquinone oxidoreductase deficiency, mitochondrial trifunctional protein deficiency, and mitochondrial short-chain 3-hydroxyacyl-CoA dehydrogenase deficiency.

In another embodiment, a vector capable of expressing EXMAD or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of EXMAD including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified EXMAD in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of EXMAD including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of EXMAD may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of EXMAD including, but not limited to, those listed above.

In a further embodiment, an antagonist of EXMAD may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of EXMAD. Examples of such disorders include, but are not limited to, those cell proliferative, immune, reproductive, neuronal, and genetic disorders described above. In one aspect, an antibody which specifically binds EXMAD may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express EXMAD.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding EXMAD may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of EXMAD including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of EXMAD may be produced using methods which are generally known in the art. In particular, purified EXMAD may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind EXMAD. Antibodies to EXMAD may also be generated using methods that are well known in the art. Such antibodies may include, but are

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with EXMAD or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Cornebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to EXMAD have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of EXMAD amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to EXMAD may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce EXMAD-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

38

complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding EXMAD, or any
 5 fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding EXMAD may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding EXMAD. Thus, complementary molecules or fragments may be used to modulate EXMAD activity, or to achieve regulation of gene function. Such technology
 10 is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding EXMAD.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to
 15 construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding EXMAD. (See, e.g., Sambrook, supra; Ausubel, 1995, supra.)

Genes encoding EXMAD can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding EXMAD. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the
 20 absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing
 25 complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding EXMAD. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may be employed. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases,
 30 transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding EXMAD.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding EXMAD. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use *in vivo*, *in vitro*, and *ex vivo*. For *ex vivo* therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such

WO 00/68380

PCT/US00/12811

therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of EXMAD, antibodies to EXMAD, and mimetics, agonists, antagonists, or inhibitors of EXMAD. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene

WO 00/68380

PCT/US00/12811

active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, or pigs.

- 5 An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

- A therapeutically effective dose refers to that amount of active ingredient, for example EXMAD or fragments thereof, antibodies of EXMAD, and agonists, antagonists or inhibitors of EXMAD, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

- 20 The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy.
- 25 Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μ g to 100,000 μ g, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art.

- 30 Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind EXMAD may be used for the

WO 00/68380

PCT/US00/12811

combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding EXMAD, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding EXMAD may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding EXMAD, or a fragment of a polynucleotide complementary to the polynucleotide encoding EXMAD, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

Methods which may also be used to quantify the expression of EXMAD include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic

variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g.,
 5 Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

In another embodiment of the invention, nucleic acid sequences encoding EXMAD may be used
 10 to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price,
 15 C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the
 20 gene encoding EXMAD on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as
 25 linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery
 30 techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or

In another embodiment of the invention, EXMAD, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between EXMAD and the agent being tested may be measured.

15 In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding EXMAD specifically compete with a test compound for binding EXMAD. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with EXMAD.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific
25 embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

30

I. Construction of cDNA Libraries

50

WO 00/68380

PCT/US00/12811

using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:26-50. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel, 1995, *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding EXMAD occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous,

WO 00/68380

PCT/US00/12811

reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories.

Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table

5 3.

V. Chromosomal Mapping of EXMAD Encoding Polynucleotides

The cDNA sequences which were used to assemble SEQ ID NO:40-50 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched
10 SEQ ID NO:40-50 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 5). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment
15 of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

The genetic map locations of SEQ ID NO:42 and SEQ ID NO:48 are described in The Invention as ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers.
20 On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters.

VI. Extension of EXMAD Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:26-50 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides
30 in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

35 High fidelity amplification was obtained by PCR using methods well known in the art. PCR

WO 00/68380

PCT/US00/12811

fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

IX. Complementary Polynucleotides

Sequences complementary to the EXMAD-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring EXMAD. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of EXMAD. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the EXMAD-encoding transcript.

X. Expression of EXMAD

Expression and purification of EXMAD is achieved using bacterial or virus-based expression systems. For expression of EXMAD in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express EXMAD upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of EXMAD in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding EXMAD by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

WO 00/68380

PCT/US00/12811

In most expression systems, EXMAD is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from EXMAD at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified EXMAD obtained by these methods can be used directly in the following activity assay.

XI. Demonstration of EXMAD Activity

An assay for EXMAD activity measures the disruption of cytoskeletal filament networks upon overexpression of EXMAD in cultured cell lines. (Reznicek, G. A. et al. (1998) J. Cell Biol. 141:209-225.) cDNA encoding EXMAD is subcloned into a mammalian expression vector that drives high levels of cDNA expression. This construct is transfected into cultured cells, such as rat kangaroo PtK2 or rat bladder carcinoma 804G cells. Actin filaments and intermediate filaments such as keratin and vimentin are visualized by immunofluorescence microscopy using antibodies and techniques well known in the art. The configuration and abundance of cytoskeletal filaments can be assessed and quantified using confocal imaging techniques. In particular, the bundling and collapse of cytoskeletal filament networks are indicative of EXMAD activity.

Alternatively, an assay for EXMAD activity measures the amount of cell aggregation induced by overexpression of EXMAD. In this assay, cultured cells such as NIH3T3 are transfected with cDNA encoding EXMAD contained within a suitable mammalian expression vector under control of a strong promoter. Cotransfection with cDNA encoding a fluorescent marker protein, such as Green Fluorescent Protein (Clontech), is useful for identifying stable transfectants. The amount of cell agglutination, or clumping, associated with transfected cells is compared with that associated with untransfected cells. The amount of cell agglutination is a direct measure of EXMAD activity.

Alternatively, cell adhesion activity in EXMAD is measured in a 96-well plate assay in which wells are first coated with EXMAD by adding solutions of EXMAD of varying concentrations to the wells. Excess EXMAD is washed off with saline, and the wells incubated with a solution of 1% bovine serum albumin to block non-specific cell binding. Aliquots of a cell suspension of a suitable cell type are then added to the wells and incubated for a period of time at 37 °C. Non-adhered cells are washed

WO 00/68380

PCT/US00/12811

populations of cells transfected with sequences encoding EXMAD and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding EXMAD and other genes of interest can be analyzed by northern analysis or microarray techniques.

XIII. Production of EXMAD Specific Antibodies

EXMAD substantially purified using polyacrylamide gel electrophoresis (PAGE: see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques. is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the EXMAD amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for anti-peptide and anti-EXMAD activity by, for example, binding the peptide or EXMAD to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIV. Purification of Naturally Occurring EXMAD Using Specific Antibodies

Naturally occurring or recombinant EXMAD is substantially purified by immunoaffinity chromatography using antibodies specific for EXMAD. An immunoaffinity column is constructed by covalently coupling anti-EXMAD antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing EXMAD are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of EXMAD (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/EXMAD binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and EXMAD is collected.

XV. Identification of Molecules Which Interact with EXMAD

EXMAD, or biologically active fragments thereof, are labeled with ^{125}I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled EXMAD, washed, and any wells with labeled EXMAD complex are assayed. Data obtained using different concentrations of EXMAD are used to calculate values for the number, affinity, and association of EXMAD with the candidate molecules.

Alternatively, molecules interacting with EXMAD are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989, *Nature* 340:245-246), or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Polyptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	26	398269	PITUNOT02	265928H1 (HNT2AGT01), 398269H1 and 398269R6 (PITUNOT02), 516201R6 (MMLR1DT01), 822473R6 (KERANOT02), 1265919F1 (BRAINOT09), 1356244F6 (LUNGNOT09), 1379344T6 (LUNGNOT10), 3586102H1 (293TF4T01), SBIA02091F1, SBIA01281F1
2	27	1258888	MENITUT03	1258888H1 (MENITUT03), 1373184H1 (BSTMNON02), 2420735R6 (SCORNON02), 2697827F3 (UTRSNOT12), 2990569T6 (KIDNFET02), SBIA02402F1, SBIA05599F1, SBIA01330F1, SBIA07058F3
3	28	1375891	LUNGNOT10	1375891H1 (LUNGNOT10), 2251462R6 (OVARUT01), 4542640H1 (THYRTMT01), SAXA00188F1, SAXA00819F1, SAXA00256F1, SAXA00101F1, SZAC00197F1
4	29	1524355	UCMCL5T01	008503T6 (HMC1NOT01), 425033R6 (BLADNOT01), 1299403T6 (BRSTNOT07), 1524355H1 (UCMCL5T01), 2480893F6 (SMCANOT01), 3072568F6 (UTRSNOR01), 3077770H1 (BONEUNT01), 3521659H1 (LUNGNOT03), 3810130H1 (CONTUT01), 4187444H1 (BRSTNOT31)
5	30	1598937	BLADNOT03	307298R6 (HEARNOT01), 637901F1 (BRSTNOT03), 872833R1 (LUNGAST01), 1360462F1 (LUNGNOT12), 1598937H1 (BLADNOT03), 1688334H1 (PROSTUT10), 2048691F6 (LIVRFET02), 3604769H1 (LUNGNOT30)
6	31	1725801	PROSNOT14	359107F1 and 359107R1 (SYNORAB01), 1725801H1 and 1725801X18C1 (PROSNOT14), 2853280H1 (CONNOT02), SBIA02129V1
7	32	1730482	BRSTTUT08	1261313R1 (SYNORAT05), 1321141F1 (BLADNOT04), 1484641F1 (CORPNOT02), 1730482H1 (BRSTTUT08), 1848053F6 (OVARNOT07), 2208990F6 (SINTFET03), 2691973F6 (LUNGNOT23), 2811183H1 (OVARNOT10), 3097712H1 (CERVNOT03), 3110665H1 (BRSTNOT17), 3738668H1 (MENTNOT01)
8	33	1810058	PROSTUT12	571697H1 (OVARNON01), 1704596F6 (DUODNOT02), 1810058H1, 1810548F6, and 1810548T6 (PROSTUT12)
9	34	2040679	HIPONON02	2040679H1 and 2040679R6 (HIPONON02), 2380160F6 (ISLTNOT01), 2621171T6 (KERANOT02), 2869976F6 (THYRNOT10)
10	35	2960051	ADRENOT09	2960051F6 and 2960051H1 (ADRENOT09), SBIA05142V1, SBIA03774V1, SBIA03935V1

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
11	36	3117318	LUNGUT013	393775H1 (TMLR2DT01), 486988H1 (HNT2AGT01), 3117318F6 and 3117318H1 (LUNGUT013), 3293662F6 (TLYJINT01), SBMA01131F1
12	37	3486992	EPIGNOT01	2615184H1 (GBLANOT01), 3486992H1 (EPIGNOT01), SBKA01303F1.comp, SBKA03723F1.comp, SBKA02206F1, SBKA01625F1.comp, SBKA02769F1, SBKA03712F1, SBKA02365F1, SBKA01975F1
13	38	4568384	HELATXT01	080350F1 (SYNORAB01), 320872H1 (EOSIHT02), 1418995F1 (KIDNNOT09), 1473647T1 (LUNGUT03), 1664971F6 (BRSTNOT09), 1738547F6 (COLNNOT22), 2367046F6 (ADRENOT07), 4568384F6 and 4568384H1 (HELATXT01)
14	39	4586187	OVARNOT13	306792F1 and 306792X11R1 (HEARNOT01), 632244F1 (KIDNNOT05), 876626R1 (LUNGAST01), 2451238F6 (ENDANOT01), 2881494F6 (UTRSTUT05), 4586187H1 (OVARNOT13), 5852878H1 (FIBAUNT02), SZZZ01051R1
15	40	401801	TMLR3DT01	401801T6 and 401801H1 (TMLR3DT01), 938106H1 (CERVNOT01), 2603123T6 (UTRSNOT10), 2607556H1 (LUNGUT07)
16	41	1721842	BLADNOT06	1721842H1, 1721842F6 and 1721842T6 (BLADNOT06), 2010387R6 (TESTNOT03), 4884119H1 (LUNLTWT01)
17	42	1833221	BRAINON01	001593H1 (U937NOT01), 389513R1 (THYMNOT02), 428370R6 (BLADNOT01), 493657H1 (HNT2NOT01), 1263824R1 (SYNORAT05), 1833221H1 (BRAINON01), 1907733F6 (CONNTUT01), 1997529R6 (BRSTTUT03), 2174658F6 (ENDCNOT03), 3114306H1 (BRSTNOT17), 3233178H1 (COLNUCT03), 4788994F6 (EPIBUNT01), 5541215H1
18	43	2041168	HIPONON02	849897R1 (NGANNOT01), 908128R2 (COLNNOT09), 999830R6 (KIDNTUT01), 1639572T6 (UTRSNOT06), 1686825F6 (PROSNOT15), 2041168H1 (HIPONON02), 2582551H1 (KIDNTUT13), 2867048H1 (KIDNNOT20), 3226063F6 (TLYJINT01), 3226063H1 (TLYJINT01), 3466031H1 (293TF2T01), 4662252H2 (BRSTTUT20), SBIA03151D1
19	44	2365794	ADRENOT07	874804H1 (LUNGAST01), 1318960T1 (BLADNOT04)

64

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
20	45	2618452	GBLANOT01	1730514F6 (BRSTTUT08), 2225286F6 (SEMVN0T01), 2225720F6 (SEMVN0T01), 2618452F6 and 2618452H1 (GBLANOT01), 2618457F6 (GBLANOT01), 3248134H1 (SEMVN0T03), 3250560H1 (SEMVN0T03), 3538176F6 (SEMVN0T04), 4068913H1 (SEMVN0T05)
21	46	2622288	KERANOT02	223636F1 (PANCNOT01), 490914R6 (HNT2AGT01), 530368R6 (BRAINOT03), 850583R1 (NGANNOT01), 898618R1 (BRSTTUT03), 932484R6 (CERVNOT01), 1302418F1 (PLACNOT02), 1368735R1 (SCORN02), 1486177F6 (CORPNOT02), 1726367F6 (PROSNOT14), 2516869H1 (LIVRTUT04), 2622288R6 and 2622288H2 (KERANOT02), 3043955H1 (HEAANOT01), 3398316H1 (UTRSNOT16), 3938796H1 (SKINBIT01), 4043471H1 (LUNGNOT35)
22	47	2806595	BLADTUT08	643445R6 (BRSTTUT02), 2806595F6 and 2806595H1 (BLADTUT08), SBRA04014D1, SBRA03510D1
23	48	2850987	BRSTTUT13	1300925F1 (BRSTNOT07), 1339833F1 (COLNTUT03), 1347463F6 (PROSNOT11), 1347463T6 (PROSNOT11), 1899642F6 (BLADTUT06), 2715093F6 (THYRN0T09), 2726463F6 (OVARUT05), 2850987H1 (BRSTTUT13), 2893008H1 (LUNGFET04), 3336701F6 (SPLNNOT10), 3341661H1 (SPLNNOT09), SXAF00652V1, SXAF03272V1
24	49	3557211	LUNGNOT31	958552H1 (KIDNNOT05), 2953281F6 and 2953281T6 (KIDNFET01), 3557211F6 and 3557211H1 (LUNGNOT31), 4306204H1 (GBLADIT01), 4420950F6 (LIVRDIT02) g2188176, g1424165
25	50	4675668	NOSEDIT02	1519431T6 (BLADTUT04), 2447058F6 (THP1NOT03), 2758306R6 (THP1AZS08), 2758306T6 (THP1AZS08), 3589494H1 (293TF5T01), 3813434H1 (TONSNOT03), 4675668H1 (NOSEDIT02), 5175727H1 (EPIBTXT01), 5313381H1 (KIDETXS02)

Table 2

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
1	309	T153 S29 S140 T153 S162 T168 S233 S258 T285 S290 T87 T159 T265	N108 N305	Signal peptide: M1-A31	similar to B. Subtilis surfactin (SFP) protein g3880360	BLAST SPSCAN
2	554	S57 S146 S265 T275 S389 T495 T496 S497 S551 S25 S34 T87 S115 S180 S212 S242 S289 T308 S361 T388 T504	N398	EGF-like domain: C98-C132 C138-C172 C178-C217 C223-C258 Cell adhesion: R363-D365 Signal peptide: M1-G21	fibulin-2 [Mus musculus] g437047	BLAST PRINTS BLOCKS PFAM MOTIFS SPSCAN HMM
3	482	S87 T37 T108 T131 S133 S148 T165 T246 S254 T256 S269 S283 S333 S404 T414 T431 S28 T29 S65 T335 T431 S446 S460 T464	N252 N445 N451	Signal peptide: M1-G22	gastric mucin [Sus scrofa] g915208	BLAST MOTIFS SPSCAN HMM
4	735	S506 S153 S243 T259 S304 T317 T378 S414 T502 S575 S670 S688 S698 S44 T116 S258 S324 S350 S356 S396 T437 T515 S610 S620 Y53	N70 N97 N144 N188 N412	Kelch motif: T284-K330 C469-G513	muskelin [Mus musculus] g3493462	BLAST PFAM

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
5	424	T209 S256 S276 T86 S311 S319 T347 S15 S354 S394 S107 Y53 S153 T217 S258 S408 T306 S358 S383		SH3 domain: V366-V422	Focal adhesion protein (FAP52) [Gallus gallus] g2217964	BLAST PFAM PRINTS BLOCKS
6	420	S293 T63 T73 S99 S101 S222 T359 T48 T63 S138 T159 S406 S409 Y53	N79 N205	Signal peptide: M1-L29 EGF-like domain: T174-C192 Cysteine-rich pattern: C181-C192	HT protein [Cricetulus griseus] g1216486	BLAST PRINTS SPSCAN MOTIFS HMM
7	795	S41 T94 S145 S243 T297 S442 S451 T687 S103 T111 T129 S184 T428 S647	N383 N387	Cell adhesion: R606-D608 von Willebrand factor type A domain: D31-L204 transmembrane domain: I50-T77	collagen type XIV [Homo sapiens] g2065167	BLAST MOTIFS PFAM PRINTS HMM
8	306	T69 T133 S255 T279 T22		Signal peptide: M1-S19 Clq domain: G149-P175 A203-I226 H227-L302	saccular collagen [Lepomis macrochirus] g687606	BLAST PFAM PRINTS BLOCKS SPSCAN HMM

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
9	338	S5 S53 S66 T119 T246 S23 T65 S102 S151 S251 T277	N217 N332	Signal peptide: M1-S22 Leucine-rich repeats domain: S102-T147 S151-I196 N197-A243	LRR47 [Drosophila melanogaster] g415947	BLAST PFAM PRINTS SPSCAN HMM
10	164	S42 S75 T160 S44 S49		Signal peptide: M1-G20 von Willebrand factor C-type domain: C103-C157	extracellular matrix protein [Homo sapiens] g3786312	BLAST PFAM BLOCKS SPSCAN HMM
11	327	S292 S30 S35 S63 T92 T14 T102 T179 S198 T285	N54 N61 N75 N85 N100 N189 N196 N213 N218 N322	Signal peptide: M1-P29 Ig domain: P81-F144 G173-A239 Transmembrane domain: V254-A276	embigin protein [Rattus norvegicus] g3355709	BLAST PFAM SPSCAN HMM
12	716	S21 T49 T54 T87 T98 S245 T315 T471 T519 T590 S624 S692 T705 S176 S384 S473 S494 T513 S542 T560 T571 T605 T613 S664 T709 Y581	N69 N96 N106 N117 N385 N517 N582 N611	Signal peptide: M1-S25 Leucine-rich repeats domain: N96-S143 N192-D239 S240-L287 R288-P337 A338-N385 Transmembrane domain: M639-F656	leucine-rich- repeat protein [Mus musculus] g1228052	BLAST PFAM PRINTS SPSCAN HMM

10009557, 061202

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
13	665	T147 S45 S86 S110 S121 T147 S160 T200 S205 S225 S247 S299 S301 S309 S335 S336 S341 S343 T386 S388 T400 T448 S506 S534 S545 S580 S581 S582 S597 S602 S615 S23 S82 S100 S162 S183 T199 S217 S221 S329 S347 T429 T501 T558 T563 T608 Y445 Y559	N119 N242 N424 N427 N634		50kDa lectin [Bombyx mori] g500858	BLAST
14	547	T60 S31 T87 T175 S213 T357 T452 T474 S476 T488 S203 T420 Y424	N15 N76 N85 N104 N128 N154 N191 N221 N242 N418	Lectin C-type domain: L473-C535 T488-L547 Cell adhesion: R256-D258	CSR1 (cellular stress response protein) [Homo sapiens] g6230372	BLAST PFAM BLOCKS MOTIFS PPROFILES SCAN
15	109	S85 S38	N22		Attractin; DPPT- L [Homo sapiens] g3676347	BLAST-GenBank MOTIFS
16	192	S10 S87 T92 T157 T165 T170 S19 S46	N8 N103	Leucine Rich Repeat Domain: L81-I94 L126-M139		BLIMPS- PRINTS MOTIFS

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
17	575	T150 S171 S299 S85 S98 S117 S118 S126 S142 S170 S203 S237 S239 S333 S415 S467 T473 S524 T557 S558 S562 S32 S92 S104 S128 S134 T149 T150 S167 S188 S260 S270 S280 S289 S389 S536	N68 N96 N234 N366 N569		axotrophin [Mus musculus] g5052031 dentin phosphoryn [Homo sapiens] g4322670	BLAST-GenBank MOTIFS
18	342	S73 S24 S82 S207 S315 S96 T176	N31 N152 N180 N193	Armadillo/beta- catenin-like repeats: A104-A113		BLIMPS-PFAM MOTIFS
19	110	S80		Signal Peptide: M1-G45 Transmembrane Domain: G48-G71 G91-Y110 Legume lectins signature: V4-F54		SPSCAN HMMER PROFILESAN MOTIFS
20	571	S482 T502 T11 T40 S88 T180 S231 T339 T383 T402 T409 T436 T447 S482 T491	N66 N229 N434 N498	Mucin domain: P101 - S430 Cystine knot domain: C481-C569	mucin [Homo sapiens] g292046	BLAST-GenBank BLAST-DOMO HMMER-PFAM MOTIFS

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
21	262	S69 S146 S172 S41 T54 T59 T101 T102 T107 Y170		Signal Peptide: M1-G25	single-pass transmembrane protein [Rattus norvegicus] g6978944	SPSCAN HMMER MOTIFS BLAST-GenBank
22	172	S29 T53 S111 S80 Y144		Signal Peptide: M1-G17 Protein proteoglycan core glycoprotein precursor cartilage repeat lectin Ig fold : G63-I149 Immunoglobulin: E52-S156	antigen [Homo sapiens] g188543 link protein [Mus musculus] g4218976	BLAST-GenBank BLAST-PRODOM BLAST-DOMO SPSCAN HMMER MOTIFS
23	571	S16 T36 T294 S396 S403 T445 S23 T176 S487	N100 N174 N434 N567	Mitochondrial energy transfer proteins signature: P404-F412 Transmembrane domains: T94-K116 F520-F539 L58-I78 I341-W362 I375-M393 I453-F472 Laminin b: Y538-K558	cell adhesion regulator [Rattus norvegicus] g4098299	BLAST- GenBank, HMMER-PFAM HMMER MOTIFS

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
24	455	S18 S23 S143 S270 S81 T186 S196 T208 S230 T240 T256 S418 S452 Y223	N138 N217 N288	Signal peptidases I signature: G43-F50 Lectin c-type: C329-S452 Cell attachment sequence: R183-D185	lectin BRA-3 [Megabalanus rosa] g407227	BLAST-GenBank BLAST-DOMO HMMER-PFAM MOTIFS
25	437	S98 T146 T160 S211 T220 T301 S55 T86 T156 S197 T369 Y265 Y334 Y350		ENP1 protein nuclear protein: E157-D431	bystin [Mus musculus] g2738509	BLAST-GenBank BLAST-PRODOM MOTIFS

Table 3

Nucleotide SEQ ID NO:	Unique Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
26	242-286	Nervous (0.264) Reproductive (0.198)	Cancer (0.462) Cell proliferation (0.242) Inflammation (0.176)	PSPORT
27	272-316	Nervous (0.438) Reproductive (0.188) Developmental (0.188)	Cancer (0.438) Cell proliferation (0.250) Inflammation (0.188)	pINCY
28	218-262	Gastrointestinal (0.244) Nervous (0.195) Reproductive (0.171)	Cancer (0.488) Inflammation (0.195) Cell proliferation (0.146)	pINCY
29	488-532 1082-1126	Reproductive (0.265) Nervous (0.206) Hematopoietic/immune (0.147)	Cancer (0.500) Cell proliferation (0.324) Inflammation (0.235)	PBLUESCRIPT
30	542-586	Reproductive (0.321) Cardiovascular (0.143) Musculoskeletal (0.143)	Cancer (0.500) Inflammation (0.107) Cell proliferation (0.107)	pINCY
31	217-261	Nervous (0.265) Reproductive (0.253) Cardiovascular (0.108)	Cancer (0.482) Inflammation (0.145) Cell proliferation (0.145)	pINCY
32	36-80	Reproductive (0.333) Gastrointestinal (0.154) Developmental (0.115)	Cancer (0.462) Inflammation (0.167) Cell proliferation (0.154)	pINCY
33	218-262	Reproductive (0.571) Gastrointestinal (0.286) Cardiovascular (0.143)	Trauma (0.286) Cancer (0.143) Inflammation (0.143)	pINCY
34	111-155	Gastrointestinal (0.364) Nervous (0.182) Cardiovascular (0.091)	Cancer (0.364) Inflammation (0.273) Cell proliferation (0.182)	PSPORT
35	271-315	Musculoskeletal (0.286) Reproductive (0.286) Cardiovascular (0.143)	Cancer (0.286) Inflammation (0.143) Neurological (0.143)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Unique Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
36	542-586 866-910	Hematopoietic/Immune (0.526) Reproductive (0.158) Nervous (0.105)	Cancer (0.368) Inflammation (0.474) Cell proliferation (0.158)	pINCY
37	811-855	Nervous (0.267) Reproductive (0.267) Musculoskeletal (0.133)	Cancer (0.600) Inflammation (0.200) Cell proliferation (0.133)	pINCY
38	380-424 974-1018	Reproductive (0.200) Gastrointestinal (0.164) Nervous (0.145)	Cancer (0.436) Cell proliferation (0.309) Inflammation (0.200)	pINCY
39	434-479 975-1019	Reproductive (0.296) Cardiovascular (0.259) Hematopoietic/Immune (0.111)	Cancer (0.315) Inflammation (0.204) Trauma (0.204)	pINCY
40	555-614	Cardiovascular (0.333) Hematopoietic/Immune (0.333) Reproductive (0.333)	Inflammation (0.667) Cancer (0.333)	PBLUESCRIPT
41	743-802	Nervous (0.353) Reproductive (0.176) Urologic (0.176)	Cancer (0.471) Inflammation (0.411) Cell Proliferation (0.118)	pINCY
42	429-488 1029-1088	Reproductive (0.213) Nervous (0.191) Cardiovascular (0.169)	Cancer (0.472) Inflammation (0.394) Cell Proliferation (0.180)	PSPORT1
43	967-1026	Nervous (0.228) Reproductive (0.213) Gastrointestinal (0.110)	Cancer (0.504) Inflammation (0.291) Cell Proliferation (0.197)	PSPORT1
44	164-223	Reproductive (0.241) Cardiovascular (0.167) Gastrointestinal (0.148)	Cancer (0.481) Inflammation (0.315) Cell Proliferation (0.167)	pINCY
45	110-169	Gastrointestinal (0.562) Reproductive (0.312) Nervous (0.062) Urologic (0.062)	Cancer (0.500) Inflammation (0.312) Cell Proliferation (0.062)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Unique Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
46	273-332	Nervous (0.347)	Cancer (0.430)	PSPORT1
	759-818	Reproductive (0.223)	Inflammation (0.364)	
		Cardiovascular (0.132)	Cell Proliferation (0.124)	
47	218-277	Gastrointestinal (0.200)	Cancer (0.533)	pINCY
		Nervous (0.200)	Inflammation (0.334)	
		Reproductive (0.200)	Cell Proliferation (0.133)	
48	341-400	Reproductive (0.294)	Cancer (0.476)	pINCY
		Gastrointestinal (0.168)	Inflammation (0.329)	
		Cardiovascular (0.126)	Cell Proliferation (0.168)	
49	266-325	Developmental (0.277)	Cell Proliferation (0.444)	pINCY
	542-601	Gastrointestinal (0.222)	Inflammation (0.444)	
		Nervous (0.167) Urologic (0.167)	Cancer (0.167)	
50	165-224	Hematopoietic/Immune (0.216)	Cancer (0.568)	pINCY
		Reproductive (0.216)	Cell Proliferation (0.324)	
		Gastrointestinal (0.135)	Inflammation (0.297)	

10005557-054202

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
26	PITUNOT02	The library was constructed using RNA obtained from Clontech. The RNA was isolated from the pituitary glands removed from a pool of 87 male and female donors, 15 to 75 years old.
27	MENITUT03	The library was constructed using RNA isolated from brain meningioma tissue removed from a 35-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated a benign neoplasm in the right cerebellopontine angle of the brain. Patient history included hypothyroidism. Family history included myocardial infarction and breast cancer.
28	LUNGNOT10	The library was constructed using RNA isolated from the lung tissue of a Caucasian male fetus, who died at 23 weeks' gestation.
29	UCMCL5T01	The UCMCL5T01 library was constructed using RNA isolated from mononuclear cells obtained from the umbilical cord blood of 12 individuals. The cells were cultured for 12 days with IL-5 before RNA was obtained from the pooled lysates.
30	BLADNOT03	The library was constructed using RNA isolated from the bladder tissue removed from an 80-year-old Caucasian female during a radical cystectomy and lymph node excision. Pathology for the associated tumor tissue indicated grade 3 invasive transitional cell carcinoma. Patient history included malignant neoplasm of the uterus, atherosclerosis, and atrial fibrillation. Family history included acute renal failure, osteoarthritis, and atherosclerosis.
31	PROSNOT14	The library was constructed using RNA isolated from diseased prostate tissue removed from a 60-year-old Caucasian male during radical prostatectomy and regional lymph node excision. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 3+4). The patient presented with elevated prostate specific antigen (PSA). Patient history included a kidney cyst and hematuria. Family history included benign hypertension, cerebrovascular disease, and arteriosclerotic coronary artery disease.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
32	BRSTTUT08	The library was constructed using RNA isolated from breast tumor tissue removed from a 45-year-old Caucasian female during unilateral extended simple mastectomy. Pathology indicated invasive nuclear grade 2-3 adenocarcinoma, ductal type, with 3 of 23 lymph nodes positive for metastatic disease. Greater than 50% of the tumor volume was in situ, both comedo and non-comedo types. Immunostains were positive for estrogen/progesterone receptors, and uninvolved tissue showed proliferative changes. The patient concurrently underwent a total abdominal hysterectomy. Patient history included valvuloplasty of mitral valve without replacement, rheumatic mitral insufficiency, and rheumatic heart disease. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.
33	PROSTUT12	The library was constructed using RNA isolated from prostate tumor tissue removed from a 65-year-old Caucasian male during a radical prostatectomy. Pathology indicated an adenocarcinoma (Gleason grade 2+2). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA).
34	HIPONON02	This normalized hippocampus library was constructed from 1.13M independent clones from a normal hippocampus library. RNA was isolated from the hippocampus tissue of a 72-year-old Caucasian female who died from an intracranial bleed. Patient history included nose cancer, hypertension, and arthritis. The normalization and hybridization conditions were adapted from Soares et al. (Proc.Natl.Acad.Sci. USA (1994) 91:9228).
35	ADREN0T09	The library was constructed using RNA isolated from left adrenal gland tissue removed from a 43-year-old Caucasian male during nephroureterectomy, regional lymph node excision, and unilateral left adrenalectomy. Pathology for the associated tumor tissue indicated a grade 2 renal cell carcinoma mass in the posterior lower pole of the left kidney with invasion into the renal pelvis.
36	LUNGUT13	The library was constructed using RNA isolated from tumorous lung tissue removed from the right upper lobe of a 47-year-old Caucasian male during a segmental lung resection. Pathology indicated invasive grade 3 (of 4) adenocarcinoma. Family history included atherosclerotic coronary artery disease, and type II diabetes.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
37	EPIGNOT01	The library was constructed using RNA isolated from epiglottic tissue removed from a 71-year-old male during laryngectomy with right parathyroid biopsy. Pathology for the associated tumor tissue indicated recurrent grade 1 papillary thyroid carcinoma.
38	HELATXT01	The library was constructed using RNA isolated from HeLa cells treated with TNF- α and IL-1 β , 10ng/nl each for 20 hours. The HeLa cell line is derived from cervical adenocarcinoma removed from a 31-year-old Black female.
39	OVARNOT13	The library was constructed using RNA isolated from left ovary tissue removed from a 47-year-old Caucasian female during a vaginal hysterectomy with bilateral salpingo-oophorectomy, and dilation and curettage. Pathology for the associated tumor tissue indicated a single intramural leiomyoma. The endometrium was in the secretory phase. The patient presented with metrorrhagia. Patient history included hyperlipidemia and benign hypertension. Family history included colon cancer, benign hypertension, atherosclerotic coronary artery disease, and breast cancer.
40	TMLR3DT01	Library was constructed using RNA isolated from non-adherent and adherent peripheral blood mononuclear cells collected from two unrelated Caucasian male donors (25 and 29 years old). Cells from each donor were purified on Ficoll Hypaque, then co-cultured for 96 hours in medium containing normal human serum at a cell density of 2×10^6 cells/ml. The non-adherent and adherent cell populations were pooled, washed once in PBS, lysed in a buffer containing GuSCN, and spun through CsCl to obtain RNA.
41	BLADNOT06	Library was constructed using RNA isolated from the posterior wall bladder tissue removed from a 66-year-old Caucasian male during a radical prostatectomy, radical cystectomy and urinary diversion. Pathology for the associated tumor tissue indicated grade 3 transitional cell carcinoma on the anterior wall of the bladder and urothelium. Patient history included lung neoplasm, and tobacco abuse in remission. Family history included a malignant breast neoplasm, tuberculosis, cerebrovascular disease, atherosclerotic coronary artery disease, and lung cancer.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
42	BRAINON01	Library was constructed and normalized from 4.88 million independent clones from a brain library. RNA was made from brain tissue removed from a 26-year-old Caucasian male during cranioplasty and excision of a cerebral meningeal lesion. Pathology for the associated tumor tissue indicated a grade 4 oligoastrocytoma in the right fronto-parietal part of the brain.
43	HIPONON02	This normalized hippocampus library was constructed from 1.13 million independent clones from a hippocampal library. RNA was isolated from the hippocampus tissue of a 72-year-old Caucasian female who died from an intracranial bleed. Patient history included nose cancer, hypertension, and arthritis. The normalization and hybridization conditions were adapted from Soares et al. (PNAS (1994) 91:9928).
44	ADRENOT07	Library was constructed using RNA isolated from adrenal tissue removed from a 61-year-old female during a bilateral adrenalectomy. Patient history included an unspecified disorder of the adrenal glands.
45	GBLANOT01	Library was constructed using RNA isolated from diseased gallbladder tissue removed from a 53-year-old Caucasian female during a cholecystectomy. Pathology indicated mild chronic cholecystitis and cholelithiasis with approximately 150 mixed gallstones. Family history included benign hypertension.
46	KERANOT02	Library was constructed using RNA isolated from epidermal breast keratinocytes (NHEK). NHEK (Clontech #CC-2501) is human breast keratinocyte cell line derived from a 30-year-old black female during breast-reduction surgery.
47	BLADTUT08	Library was constructed using RNA isolated from bladder tumor tissue removed from a 72-year-old Caucasian male during a radical cystectomy and prostatectomy. Pathology indicated an invasive grade 3 (of 3) transitional cell carcinoma in the right bladder base. Patient history included pure hypercholesterolemia and tobacco abuse. Family history included myocardial infarction, cerebrovascular disease, brain cancer, and myocardial infarction.
48	BRSTTUT13	Library was constructed using RNA isolated from breast tumor tissue removed from the right breast of a 46-year-old Caucasian female during a unilateral extended simple mastectomy with breast reconstruction. Pathology indicated an invasive grade 3 adenocarcinoma, ductal type with apocrine features and greater than 50% intraductal component. Patient history included breast cancer.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
49	LUNGNOT31	Library was constructed using RNA isolated from right middle lobe lung tissue removed from a 63-year-old Caucasian male. Pathology for the associated tumor indicated grade 3 adenocarcinoma. Patient history included an abdominal aortic aneurysm, cardiac dysrhythmia, atherosclerotic coronary artery disease, hiatal hernia, chronic sinusitis, and lupus. Family history included acute myocardial infarction and atherosclerotic coronary artery disease.
50	NOSEDIT02	The library was constructed using RNA isolated from nasal polyp tissue.

10009557 061204

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Normalized quality score > GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <i>supra</i> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

WO 00/68380

PCT/US00/12811

10. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:

- a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50,
- b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50,
- c) a polynucleotide sequence complementary to a),
- d) a polynucleotide sequence complementary to b), and
- e) an RNA equivalent of a)-d).

11. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 10.

12. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 10, the method comprising:

- a) hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

13. A method of claim 12, wherein the probe comprises at least 30 contiguous nucleotides.

14. A method of claim 12, wherein the probe comprises at least 60 contiguous nucleotides.

15. A pharmaceutical composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

16. A method for treating a disease or condition associated with decreased expression of functional EXMAD, comprising administering to a patient in need of such treatment the pharmaceutical composition of claim 15.

17. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and

Replaced by
Dr. Talebi

100094557 .06.1202

WO 00/68380

PCT/US00/12811

b) detecting agonist activity in the sample.

18. A pharmaceutical composition comprising an agonist compound identified by a method of claim 17 and a pharmaceutically acceptable excipient.

5

19. A method for treating a disease or condition associated with decreased expression of functional EXMAD, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 18.

10 20. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

15 21. A pharmaceutical composition comprising an antagonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

20 22. A method for treating a disease or condition associated with overexpression of functional EXMAD, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 21.

23. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 4, the method comprising:

- 25
- a) exposing a sample comprising the target polynucleotide to a compound, and
 - b) detecting altered expression of the target polynucleotide.

b) detecting agonist activity in the sample.

18. A pharmaceutical composition comprising an agonist compound identified by a method of claim 17 and a pharmaceutically acceptable excipient.

5

19. A method for treating a disease or condition associated with decreased expression of functional EXMAD, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 18.

10

20. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

15

21. A pharmaceutical composition comprising an antagonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

22. A method for treating a disease or condition associated with overexpression of functional EXMAD, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 21.

20

23. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 4, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

25

30

35

00 / 12811
14 AUG 2001

- 35

PF-0693 PCT

5

10

15

20

25

30

35

35. A method of diagnosing a condition or disease associated with the expression of EXMAD in a subject, comprising administering to said subject an effective amount of the composition of claim 34.

5 36. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 9, the method comprising:

- a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 1-25, or an immunogenic fragment thereof, under conditions to elicit an antibody response,
- 10 b) isolating antibodies from said animal, and
- c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 1-25.

15 37. An antibody produced by a method of claim 36.

38. A composition comprising the antibody of claim 37 and a suitable carrier.

20 39. A method of making a monoclonal antibody with the specificity of the antibody of claim 9, the method comprising:

- a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 1-25, or an immunogenic fragment thereof, under conditions to elicit an antibody response,
- b) isolating antibody producing cells from the animal,
- 25 c) fusing the antibody producing cells with immortalized cells to form monoclonal antibody-producing hybridoma cells,
- d) culturing the hybridoma cells, and
- e) isolating from the culture monoclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of
- 30 SEQ ID NO: 1-25.

40. A monoclonal antibody produced by a method of claim 39.

41. A composition comprising the antibody of claim 40 and a suitable carrier.

42. The antibody of claim 9, wherein the antibody is produced by screening a Fab expression library.

43. The antibody of claim 9, wherein the antibody is produced by screening a recombinant immunoglobulin library.

44. A method of detecting a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-25 in a sample, the method comprising:

- a) incubating the antibody of claim 9 with a sample under conditions to allow specific binding of the antibody and the polypeptide, and
- b) detecting specific binding, wherein specific binding indicates the presence of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-25 in the sample.

45. A method of purifying a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-25 from a sample, the method comprising:

- a) incubating the antibody of claim 9 with a sample under conditions to allow specific binding of the antibody and the polypeptide, and
- b) separating the antibody from the sample and obtaining the purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-25.

46. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

47. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.

48. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.

49. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:4.

50. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:5.

51. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID

NO:6.

69. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:24.

70. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID

71. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID

72. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID

73. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID

74. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID

75. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID

76. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID

77. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID

78. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID

79. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID

91. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID
35 NO:46.

PF-0693 PCT

PCT/US 00/12811
IPEA/US 14 AUG 2001

92. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID
NO:47.

93. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID
5 NO:48.

94. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID
NO:49.

10 95. A polynucleotide of claim 10, comprising the polynucleotide sequence of SEQ ID
NO:50.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 November 2000 (16.11.2000)

PCT

(10) International Publication Number
WO 00/68380 A3

(51) International Patent Classification⁷: C12N 15/12,
C07K 14/78, 14/47, C12N 15/63, A01K 67/027, C07K
16/18, C12Q 1/68, A61K 38/17, G01N 33/68

(21) International Application Number: PCT/US00/12811

(22) International Filing Date: 10 May 2000 (10.05.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/133,643 11 May 1999 (11.05.1999) US
60/150,409 23 August 1999 (23.08.1999) US

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier applications:

US 60/133,643 (CIP)
Filed on 11 May 1999 (11.05.1999)
US 60/150,409 (CIP)
Filed on 23 August 1999 (23.08.1999)

(71) Applicant (for all designated States except US): INCYTE
GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo
Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BANDMAN, Olga
[US/US]; 366 Anna Avenue, Mountain View, CA 94043
(US). HILLMAN, Jennifer, L. [US/US]; 230 Monroe
Drive, #12, Mountain View, CA 94040 (US). TANG,
Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA
95118 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa

Clara, CA 95054 (US). YUE, Henry [US/US]; 826 Lois
Avenue, Sunnyvale, CA 94086 (US). BAUGHN, Mariah,
R. [US/US]; 14244 Santiago Road, San Leandro, CA
94577 (US). LU, Dyung, Aina, M. [US/US]; 55 Park
Belmont Place, San Jose, CA 95136 (US). AZIMZAI,
Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA
94545 (US).

(74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics,
Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,
ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,
SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ,
VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

(88) Date of publication of the international search report:
19 April 2001

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: EXTRACELLULAR MATRIX AND ADHESION-ASSOCIATED PROTEINS

(57) Abstract: The invention provides human extracellular matrix and adhesion-associated proteins (EXMAD) and polynucleotides which identify and encode EXMAD. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of EXMAD.

WO 00/68380 A3

Docket No.: PF-0693 USN

Country	Number	Filing Date	Priority Claimed
_____	_____	_____	// Yes // No
_____	_____	_____	// Yes // No

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

Application		Status (Pending,
Serial No.	Filed	Abandoned, Patented)
60/133,643	May 11, 1999	Expired
60/150,409	August 23, 1999	Expired

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in said prior application(s) in the manner required by the first paragraph of Title 35, United States Code §112, I acknowledge my duty to disclose material information as defined in Title 37 Code of Federal Regulations, §1.56(a) which occurred between the filing date(s) of the prior application(s) and the national or Patent Cooperation Treaty international filing date of this application:

Application		Status (Pending,
Serial No.	Filed	Abandoned, Patented)
_____	_____	_____

I hereby appoint the following:

Lucy J. Billings	Reg. No. <u>36,749</u>
Michael C. Cerrone	Reg. No. <u>39,132</u>
Diana Hamlet-Cox	Reg. No. <u>33,302</u>
Richard C. Ekstrom	Reg. No. <u>37,027</u>
Barrie D. Greene	Reg. No. <u>46,740</u>
Matthew R. Kaser	Reg. No. <u>44,817</u>
Lynn E. Murry	Reg. No. <u>42,918</u>
Shirley A. Recipon	Reg. No. <u>47,016</u>
Susan K. Sather	Reg. No. <u>44,316</u>
Michelle M. Stempien	Reg. No. <u>41,327</u>
David G. Streeter	Reg. No. <u>43,168</u>
Stephen Todd	Reg. No. <u>47,139</u>
P. Ben Wang	Reg. No. <u>41,420</u>

13

respectively and individually, as my patent attorneys and/or agents, with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith. Please address all communications to:

Docket No.: PF-0693 USN

LEGAL DEPARTMENT
INCYTE GENOMICS, INC.
3160 PORTER DRIVE, PALO ALTO, CA 94304

TEL: 650-855-0555 FAX: 650-849-8886 or 650-845-4166

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

First Joint Inventor:

1-00
Full name: Olga Bandman
Signature: Olga Bandman
Date: 26 September, 2001
Citizenship: United States
Residence: Mountain View, California CA
P.O. Address: 366 Anna Avenue
Mountain View, California 94043

Second Joint Inventor:

2-00
Full name: Jennifer L. Hillman
Signature: Jennifer L. Hillman
Date: Sept 21, 2001
Citizenship: United States
Residence: Mountain View, California CA
P.O. Address: 230 Monroe Drive, #17
Mountain View, California 94040

Docket No.: PF-0693 USN

Third Joint Inventor:

Full name: 300 Y. Tom Tang
Signature: [Signature]
Date: Sept -10, 2001
Citizenship: United States
Residence: San Jose, California CA
P.O. Address: 4230 Ranwick Court
San Jose, California 95118

Fourth Joint Inventor:

Full name: 400 Preeti Lal
Signature: [Signature]
Date: September 10, 2001
Citizenship: India
Residence: Santa Clara, California CA
P.O. Address: P.O. Box 5142
Santa Clara, California 95056

Fifth Joint Inventor:

Full name: 500 Henry Yue
Signature: [Signature]
Date: September 24, 2001
Citizenship: United States
Residence: Sunnyvale, California CA
P.O. Address: 826 Lois Avenue
Sunnyvale, California 94087

Docket No.: PF-0693 USN

Sixth Joint Inventor:

Full name: 6-00 Mariah R. Baughn
 Signature: Mariah R. Baughn
 Date: September 5, 2001
 Citizenship: United States
 Residence: San Leandro, California CA
 P.O. Address: 14244 Santiago Road
San Leandro, California 94577

Seventh Joint Inventor:

Full name: 7-00 Dyung Aina M. Lu
 Signature: Aina Lu
 Date: September 7, 2001
 Citizenship: United States
 Residence: San Jose, California CA
 P.O. Address: 233 Coy Drive
San Jose, California 95123

Eighth Joint Inventor:

Full name: 8-00 Yalda Azimzai
 Signature: Yalda Azimzai
 Date: September 13, 2001
 Citizenship: United States
 Residence: Castro Valley, California CA
 P.O. Address: 5518 Boulder Canyon Drive
Castro Valley, California 94552

WO 00/68380

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.
 BANDMAN, Olga
 HILLMAN, Jennifer L.
 TANG, Y. Tom
 LAL, Preeti
 YUE, Henry
 BAUGHN, Mariah R.
 LU, Dyung Aina M.
 AZIMZAI, Yalda

<120> EXTRACELLULAR MATRIX AND ADHESION-ASSOCIATED PROTEINS

<130> PF-0693 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/133,643; 60/150,409

<151> 1999-05-11; 1999-08-23

<160> 50

<170> PERL Program

<210> 1

<211> 309

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 398269CD1

<400> 1

Met	Val	Phe	Pro	Ala	Lys	Arg	Phe	Cys	Leu	Val	Pro	Ser	Met	Glu
1				5					10					15
Gly	Val	Arg	Trp	Ala	Phe	Ser	Cys	Gly	Thr	Trp	Leu	Pro	Ser	Arg
				20					25					30
Ala	Glu	Trp	Leu	Leu	Ala	Val	Arg	Ser	Ile	Gln	Pro	Glu	Glu	Lys
				35					40					45
Glu	Arg	Ile	Gly	Gln	Phe	Val	Phe	Ala	Arg	Asp	Ala	Lys	Ala	Ala
				50					55					60
Met	Ala	Gly	Arg	Leu	Met	Ile	Arg	Lys	Leu	Val	Ala	Glu	Lys	Leu
				65					70					75
Asn	Ile	Pro	Trp	Asn	His	Ile	Arg	Leu	Gln	Arg	Thr	Ala	Lys	Gly
				80					85					90
Lys	Pro	Val	Leu	Ala	Lys	Asp	Ser	Ser	Asn	Pro	Tyr	Pro	Asn	Phe
				95					100					105
Asn	Phe	Asn	Ile	Ser	His	Gln	Gly	Asp	Tyr	Ala	Val	Leu	Ala	Ala
				110					115					120
Glu	Pro	Glu	Leu	Gln	Val	Gly	Ile	Asp	Ile	Met	Lys	Thr	Ser	Phe
				125					130					135
Pro	Gly	Arg	Gly	Ser	Ile	Pro	Glu	Phe	Phe	His	Ile	Met	Lys	Arg
				140					145					150
Lys	Phe	Thr	Asn	Lys	Glu	Trp	Glu	Thr	Ile	Arg	Ser	Phe	Lys	Asp
				155					160					165
Glu	Trp	Thr	Gln	Leu	Asp	Met	Phe	Tyr	Arg	Asn	Trp	Ala	Leu	Lys
				170					175					180
Glu	Ser	Phe	Ile	Lys	Ala	Ile	Gly	Val	Gly	Leu	Gly	Phe	Glu	Leu
				185					190					195
Gln	Arg	Leu	Glu	Phe	Asp	Leu	Ser	Pro	Leu	Asn	Leu	Asp	Ile	Gly
				200					205					210
Gln	Val	Tyr	Lys	Glu	Thr	Arg	Leu	Phe	Leu	Asp	Gly	Glu	Glu	Glu
				215					220					225
Lys	Glu	Trp	Ala	Phe	Glu	Glu	Ser	Lys	Ile	Asp	Glu	His	His	Phe

WO 00/68380

PCT/US00/12811

```

230                235                240
Val Ala Val Ala Leu Arg Lys Pro Asp Gly Ser Arg His Gln Asp
245                250                255
Val Pro Ser Gln Asp Asp Ser Lys Pro Thr Gln Arg Gln Phe Thr
260                265                270
Ile Leu Asn Phe Asn Asp Leu Met Ser Ser Ala Val Pro Met Thr
275                280                285
Pro Glu Asp Pro Ser Phe Trp Asp Cys Phe Cys Phe Thr Glu Glu
290                295                300
Ile Pro Ile Arg Asn Gly Thr Lys Ser
305

<210> 2
<211> 554
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1258888CD1

<400> 2
Met Pro Leu Pro Trp Ser Leu Ala Leu Pro Leu Leu Leu Ser Trp
1      5      10      15
Val Ala Gly Gly Phe Gly Asn Ala Ala Ser Ala Arg His His Gly
20      25      30
Leu Leu Ala Ser Ala Arg Gln Pro Gly Val Cys His Tyr Gly Thr
35      40      45
Lys Leu Ala Cys Cys Tyr Gly Trp Arg Arg Asn Ser Lys Gly Val
50      55      60
Cys Glu Ala Thr Cys Glu Pro Gly Cys Lys Phe Gly Glu Cys Val
65      70      75
Gly Pro Asn Lys Cys Arg Cys Phe Pro Gly Tyr Thr Gly Lys Thr
80      85      90
Cys Ser Gln Asp Val Asn Glu Cys Gly Met Lys Pro Arg Pro Cys
95      100     105
Gln His Arg Cys Val Asn Thr His Gly Ser Tyr Lys Cys Phe Cys
110     115     120
Leu Ser Gly His Met Leu Met Pro Asp Ala Thr Cys Val Asn Ser
125     130     135
Arg Thr Cys Ala Met Ile Asn Cys Gln Tyr Ser Cys Glu Asp Thr
140     145     150
Glu Glu Gly Pro Gln Cys Leu Cys Pro Ser Ser Gly Leu Arg Leu
155     160     165
Ala Pro Asn Gly Arg Asp Cys Leu Asp Ile Asp Glu Cys Ala Ser
170     175     180
Gly Lys Val Ile Cys Pro Tyr Asn Arg Arg Cys Val Asn Thr Phe
185     190     195
Gly Ser Tyr Tyr Cys Lys Cys His Ile Gly Phe Glu Leu Gln Tyr
200     205     210
Ile Ser Gly Arg Tyr Asp Cys Ile Asp Ile Asn Glu Cys Thr Met
215     220     225
Asp Ser His Thr Cys Ser His His Ala Asn Cys Phe Asn Thr Gln
230     235     240
Gly Ser Phe Lys Cys Lys Cys Lys Gln Gly Tyr Lys Gly Asn Gly
245     250     255
Leu Arg Cys Ser Ala Ile Pro Glu Asn Ser Val Lys Glu Val Leu
260     265     270
Arg Ala Pro Gly Thr Ile Lys Asp Arg Ile Lys Lys Leu Leu Ala
275     280     285
His Lys Asn Ser Met Lys Lys Lys Ala Lys Ile Lys Asn Val Thr
290     295     300
Pro Glu Pro Thr Arg Thr Pro Thr Pro Lys Val Asn Leu Gln Pro
305     310     315
Phe Asn Tyr Glu Glu Ile Val Ser Arg Gly Gly Asn Ser His Gly
320     325     330
Gly Lys Lys Gly Asn Glu Glu Lys Met Lys Glu Gly Leu Glu Asp
335     340     345

```


WO 00/68380

PCT/US00/12811

Thr	Pro	Ser	Trp	215	Ser	Pro	Gly	Ser	Asp	220	Val	Thr	Leu	Leu	Ala	225	Glu
				230	Val	Thr	Asn	Ile	Glu	235	Val	Ile	Asn	Cys	Ser	240	Ile
Ala	Leu	Val	Thr	245	Thr	Thr	Ser	Ser		250	Pro	Gly	Ala	Ser		255	Asp
Thr	Glu	Ile	Glu	260	Pro	Thr	Glu	Gly	Val	265	Lys	Ala	Ser	Ser	Thr	270	Ser
Ile	Asp	Leu	Ile	275	Leu	Pro	Asp	Ser	Thr	280	Glu	Ala	Lys	Pro	His	285	Ile
Asp	Pro	Pro	Ala	290	Ala	Ser	Ala	Glu	Thr	295	Leu	Ser	Thr	Ala	Gly	300	Thr
Thr	Glu	Val	Thr	305	Ala	Pro	His	Ala	Thr	310	Val	Gly	Thr	Pro	Leu	315	Pro
Thr	Glu	Ser	Ala	320	Thr	Glu	Arg	Glu	Val	325	Thr	Ala	Pro	Gly	Ala	330	Thr
Thr	Asn	Ser	Ala	335	Ala	Leu	Val	Thr	Val	340	Ser	Arg	Asn	Pro	Leu	345	Glu
Thr	Leu	Ser	Gly	350	Leu	Ser	Val	Glu	Thr	355	Pro	Ser	Tyr	Val	Lys	360	Val
Glu	Thr	Ser	Ala	365	Pro	Val	Ser	Ile	Glu	370	Ala	Gly	Ser	Ala	Val	375	Gly
Ser	Gly	Ala	Ala	380	Ala	Gly	Ser	Ser		385	Ala	Ser	Ser	Tyr	Ser	390	Pro
Lys	Thr	Thr	Ser	395	Leu	Lys	Asn	Phe	Thr	400	Pro	Ser	Glu	Thr	Pro	405	Thr
Ser	Glu	Ala	Ala	410	Thr	Lys	Gly	Pro	Phe	415	Pro	Thr	Ser	Arg	Asp	420	Pro
Met	Asp	Ile	Ala	425	Pro	Pro	Thr	Thr	Thr	430	Asn	Ser	Ser	Arg	Gly	435	Thr
Leu	Pro	Ser	Val	440	Ala	Lys	Ile	Thr	Thr	445	Ser	Ala	Lys	Thr	Thr	450	Met
Asn	Ser	Thr	Leu	455	Pro	Arg	Pro	Arg	Leu	460	Pro	Gly	Arg	Gly	Arg	465	Pro
Lys	Pro	Gln	Gln	470						475						480	
Gln	Thr																

<210> 4

<211> 735

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1524355CD1

<400> 4

Met	Ala	Ala	Gly	Gly	Ala	Val	Ala	Ala	Ala	Pro	Glu	Cys	Arg	Leu			
1				5					10					15			
Leu	Pro	Tyr	Ala	Leu	His	Lys	Trp	Ser	Ser	Phe	Ser	Ser	Thr	Tyr			
				20					25					30			
Leu	Pro	Glu	Asn	Ile	Leu	Val	Asp	Lys	Pro	Asn	Asp	Gln	Ser	Ser			
				35					40					45			
Arg	Trp	Ser	Ser	Glu	Ser	Asn	Tyr	Pro	Pro	Gln	Tyr	Leu	Ile	Leu			
				50					55					60			
Lys	Leu	Glu	Arg	Pro	Ala	Ile	Val	Gln	Asn	Ile	Thr	Phe	Gly	Lys			
				65					70					75			
Tyr	Glu	Lys	Thr	His	Val	Cys	Asn	Leu	Lys	Lys	Phe	Lys	Val	Phe			
				80					85					90			
Gly	Gly	Met	Asn	Glu	Glu	Asn	Met	Thr	Glu	Leu	Leu	Ser	Ser	Gly			
				95					100					105			
Leu	Lys	Asn	Asp	Tyr	Asn	Lys	Glu	Thr	Phe	Thr	Leu	Lys	His	Lys			
				110					115					120			
Ile	Asp	Glu	Gln	Met	Phe	Pro	Cys	Arg	Phe	Ile	Lys	Ile	Val	Pro			
				125					130					135			
Leu	Leu	Ser	Trp	Gly	Pro	Ser	Phe	Asn	Phe	Ser	Ile	Trp	Tyr	Val			
				140					145					150			

WO 00/68380

PCT/US00/12811

Glu	Leu	Ser	Gly	Ile	Asp	Asp	Pro	Asp	Ile	Val	Gln	Pro	Cys	Leu	155
Asn	Trp	Tyr	Ser	Lys	Tyr	Arg	Glu	Gln	Glu	Ala	Ile	Arg	Leu	Cys	160
Leu	Lys	His	Phe	Arg	Gln	His	Asn	Tyr	Thr	Glu	Ala	Phe	Glu	Ser	170
Leu	Gln	Lys	Lys	Thr	Lys	Ile	Ala	Leu	Glu	His	Pro	Met	Leu	Thr	185
Asp	Ile	His	Asp	Lys	Leu	Val	Leu	Lys	Gly	Asp	Phe	Asp	Ala	Cys	190
Glu	Glu	Leu	Ile	Glu	Lys	Ala	Val	Asn	Asp	Gly	Leu	Phe	Asn	Gln	200
Tyr	Ile	Ser	Gln	Gln	Glu	Tyr	Lys	Pro	Arg	Trp	Ser	Gln	Ile	Ile	215
Pro	Lys	Ser	Thr	Lys	Gly	Asp	Gly	Glu	Asp	Asn	Arg	Pro	Gly	Met	220
Arg	Gly	Gly	His	Gln	Met	Val	Ile	Asp	Val	Gln	Thr	Glu	Thr	Val	230
Tyr	Leu	Phe	Gly	Gly	Trp	Asp	Gly	Thr	Gln	Asp	Leu	Ala	Asp	Phe	245
Trp	Ala	Tyr	Ser	Val	Lys	Glu	Asn	Gln	Trp	Thr	Cys	Ile	Ser	Arg	250
Asp	Thr	Glu	Lys	Glu	Asn	Gly	Pro	Ser	Ala	Arg	Ser	Cys	His	Lys	260
Met	Cys	Ile	Asp	Ile	Gln	Arg	Arg	Gln	Ile	Tyr	Thr	Leu	Gly	Arg	275
Tyr	Leu	Asp	Ser	Ser	Val	Arg	Asn	Ser	Lys	Ser	Leu	Lys	Ser	Asp	285
Phe	Tyr	Arg	Tyr	Asp	Ile	Asp	Thr	Asn	Thr	Trp	Met	Leu	Leu	Ser	290
Glu	Asp	Thr	Ala	Ala	Asp	Gly	Gly	Pro	Lys	Leu	Val	Phe	Asp	His	305
Gln	Met	Cys	Met	Asp	Ser	Glu	Lys	His	Met	Ile	Tyr	Thr	Phe	Gly	310
Gly	Arg	Ile	Leu	Thr	Cys	Asn	Gly	Ser	Val	Asp	Asp	Ser	Arg	Ala	320
Ser	Glu	Pro	Gln	Phe	Ser	Gly	Leu	Phe	Ala	Phe	Asn	Cys	Gln	Cys	335
Gln	Thr	Trp	Lys	Leu	Leu	Arg	Glu	Asp	Ser	Cys	Asn	Ala	Gly	Pro	340
Glu	Asp	Ile	Gln	Ser	Arg	Ile	Gly	His	Cys	Met	Leu	Phe	His	Ser	350
Lys	Asn	Arg	Cys	Leu	Tyr	Val	Phe	Gly	Gly	Gln	Arg	Ser	Lys	Thr	365
Tyr	Leu	Asn	Asp	Phe	Phe	Ser	Tyr	Asp	Val	Asp	Ser	Asp	His	Val	370
Asp	Ile	Ile	Ser	Asp	Gly	Thr	Lys	Lys	Asp	Ser	Gly	Met	Val	Pro	385
Met	Thr	Gly	Phe	Thr	Gln	Arg	Ala	Thr	Ile	Asp	Pro	Glu	Leu	Asn	395
Glu	Ile	His	Val	Leu	Ser	Gly	Leu	Ser	Lys	Asp	Lys	Glu	Lys	Arg	400
Glu	Glu	Asn	Val	Arg	Asn	Ser	Phe	Trp	Ile	Tyr	Asp	Ile	Val	Arg	410
Asn	Ser	Trp	Ser	Cys	Val	Tyr	Lys	Asn	Asp	Gln	Ala	Ala	Lys	Asp	425
Asn	Pro	Thr	Lys	Ser	Leu	Gln	Glu	Glu	Glu	Pro	Cys	Pro	Arg	Phe	430
Ala	His	Gln	Leu	Val	Tyr	Asp	Glu	Leu	His	Lys	Val	His	Tyr	Leu	440
Phe	Gly	Gly	Asn	Pro	Gly	Lys	Ser	Cys	Ser	Pro	Lys	Met	Arg	Leu	455
Asp	Asp	Phe	Trp	Ser	Leu	Lys	Leu	Cys	Arg	Pro	Ser	Lys	Asp	Tyr	460
Leu	Leu	Arg	His	Cys	Lys	Tyr	Leu	Ile	Arg	Lys	His	Arg	Phe	Glu	475
Glu	Lys	Ala	Gln	Val	Asp	Pro	Leu	Ser	Ala	Leu	Lys	Tyr	Leu	Gln	485

WO 00/68380

PCT/US00/12811

```

        650                      655                      660
Asn Asp Leu Tyr Ile Thr Val Asp His Ser Asp Pro Glu Glu Thr
        665                      670                      675
Lys Glu Phe Gln Leu Leu Ala Ser Ala Leu Phe Lys Ser Gly Ser
        680                      685                      690
Asp Phe Thr Ala Leu Gly Phe Ser Asp Val Asp His Thr Tyr Ala
        695                      700                      705
Gln Arg Thr Gln Leu Phe Asp Thr Leu Val Asn Phe Phe Pro Asp
        710                      715                      720
Ser Met Thr Pro Pro Lys Gly Asn Leu Val Asp Leu Ile Thr Leu
        725                      730                      735
<210> 5
<211> 424
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1598937CD1

<400> 5
Met Ala Pro Glu Glu Asp Ala Gly Gly Glu Ala Leu Gly Gly Ser
  1          5          10          15
Phe Trp Glu Ala Gly Asn Tyr Arg Arg Thr Val Gln Arg Val Glu
  20          25          30
Asp Gly His Arg Leu Cys Gly Asp Leu Val Ser Cys Phe Gln Glu
  35          40          45
Arg Ala Arg Ile Glu Lys Ala Tyr Ala Gln Gln Leu Ala Asp Trp
  50          55          60
Ala Arg Lys Trp Arg Gly Thr Val Glu Lys Gly Pro Gln Tyr Gly
  65          70          75
Thr Leu Glu Lys Ala Trp His Ala Phe Phe Thr Ala Ala Glu Arg
  80          85          90
Leu Ser Ala Leu His Leu Glu Val Arg Glu Lys Leu Gln Gly Gln
  95          100         105
Asp Ser Glu Arg Val Arg Ala Trp Gln Arg Gly Ala Phe His Arg
  110         115         120
Pro Val Leu Gly Gly Phe Arg Glu Ser Arg Ala Ala Glu Asp Gly
  125         130         135
Phe Arg Lys Ala Gln Lys Pro Trp Leu Lys Arg Leu Lys Glu Val
  140         145         150
Glu Ala Ser Lys Lys Ser Tyr His Ala Ala Arg Lys Asp Glu Lys
  155         160         165
Thr Ala Gln Thr Arg Glu Ser His Ala Lys Ala Asp Ser Ala Val
  170         175         180
Ser Gln Glu Gln Leu Arg Lys Leu Gln Glu Arg Val Glu Arg Cys
  185         190         195
Ala Lys Glu Ala Glu Lys Thr Lys Ala Gln Tyr Glu Gln Thr Leu
  200         205         210
Ala Glu Leu His Arg Tyr Thr Pro Arg Tyr Met Glu Asp Met Glu
  215         220         225
Gln Ala Phe Glu Thr Cys Gln Ala Ala Glu Arg Gln Arg Leu Leu
  230         235         240
Phe Phe Lys Asp Met Leu Leu Thr Leu His Gln His Leu Asp Leu
  245         250         255
Ser Ser Ser Glu Lys Phe His Glu Leu His Arg Asp Leu His Gln
  260         265         270
Gly Ile Glu Ala Ala Ser Asp Glu Glu Asp Leu Arg Trp Trp Arg
  275         280         285
Ser Thr His Gly Pro Gly Met Ala Met Asn Trp Pro Gln Phe Glu
  290         295         300
Glu Trp Ser Leu Asp Thr Gln Arg Thr Ile Ser Arg Lys Glu Lys
  305         310         315
Gly Gly Arg Ser Pro Asp Glu Val Thr Leu Thr Ser Ile Val Pro
  320         325         330
Thr Arg Asp Gly Thr Ala Pro Pro Pro Gln Ser Pro Gly Ser Pro
  335         340         345

```

WO 00/68380

PCT/US00/12811

Gly Thr Gly Gln Asp	Glu Glu Trp Ser Asp	Glu Glu Ser Pro Arg
350	355	360
Lys Ala Ala Thr Gly	Val Arg Val Arg Ala	Leu Tyr Asp Tyr Ala
365	370	375
Gly Gln Glu Ala Asp	Glu Leu Ser Phe Arg	Ala Gly Glu Glu Leu
380	385	390
Leu Lys Met Ser Glu	Glu Asp Glu Gln Gly	Trp Cys Gln Gly Gln
395	400	405
Leu Gln Ser Gly Arg	Ile Gly Leu Tyr Pro	Ala Asn Tyr Val Glu
410	415	420
Cys Val Gly Ala		

<210> 6
 <211> 420
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1725801CD1

<400> 6

Met Ala Pro Trp Pro	Pro Lys Gly Leu Val	Pro Ala Val Leu Trp
1	5	10
Gly Leu Ser Leu Phe	Leu Asn Leu Pro Gly	Pro Ile Trp Leu Gln
20	25	30
Pro Ser Pro Pro Pro	Gln Ser Ser Pro Pro	Pro Gln Pro His Pro
35	40	45
Cys His Thr Cys Arg	Gly Leu Val Asp Ser	Phe Asn Lys Gly Leu
50	55	60
Glu Arg Thr Ile Arg	Asp Asn Phe Gly Gly	Gly Asn Thr Ala Trp
65	70	75
Glu Glu Glu Asn Leu	Ser Lys Tyr Lys Asp	Ser Glu Thr Arg Leu
80	85	90
Val Glu Val Leu Glu	Gly Val Cys Ser Lys	Ser Asp Phe Glu Cys
95	100	105
His Arg Leu Leu Glu	Leu Ser Glu Glu Leu	Val Glu Ser Trp Trp
110	115	120
Phe His Lys Gln Gln	Glu Ala Pro Asp Leu	Phe Gln Trp Leu Cys
125	130	135
Ser Asp Ser Leu Lys	Leu Cys Cys Pro Ala	Gly Thr Phe Gly Pro
140	145	150
Ser Cys Leu Pro Cys	Pro Gly Gly Thr Glu	Arg Pro Cys Gly Gly
155	160	165
Tyr Gly Gln Cys Glu	Gly Glu Gly Thr Arg	Gly Gly Ser Gly His
170	175	180
Cys Asp Cys Gln Ala	Gly Tyr Gly Gly Glu	Ala Cys Gly Gln Cys
185	190	195
Gly Leu Gly Tyr Phe	Glu Ala Glu Arg Asn	Ala Ser His Leu Val
200	205	210
Cys Ser Ala Cys Phe	Gly Pro Cys Ala Arg	Cys Ser Gly Pro Glu
215	220	225
Glu Ser Asn Cys Leu	Gln Cys Lys Lys Gly	Trp Ala Leu His His
230	235	240
Leu Lys Cys Val Asp	Ile Asp Glu Cys Gly	Thr Glu Gly Ala Asn
245	250	255
Cys Gly Ala Asp Gln	Phe Cys Val Asn Thr	Glu Gly Ser Tyr Glu
260	265	270
Cys Arg Asp Cys Ala	Lys Ala Cys Leu Gly	Cys Met Gly Ala Gly
275	280	285
Pro Gly Arg Cys Lys	Lys Cys Ser Pro Gly	Tyr Gln Gln Val Gly
290	295	300
Ser Lys Cys Leu Asp	Val Asp Glu Cys Glu	Thr Glu Val Cys Pro
305	310	315
Gly Glu Asn Lys Gln	Cys Glu Asn Thr Glu	Gly Gly Tyr Arg Cys
320	325	330
Ile Cys Ala Glu Gly	Tyr Lys Gln Met Glu	Gly Ile Cys Val Lys


```
<220>  
<221> misc_feature
```


WO 00/68380

PCT/US00/12811

Val	Arg	Val	Asp	Met	Arg	Met	Leu	Cys	Leu	Lys	Ser	Leu	Arg	Lys
				95					100					105
Leu	Asp	Leu	Ser	His	Asn	His	Ile	Lys	Lys	Leu	Pro	Ala	Thr	Ile
				110					115					120
Gly	Asp	Leu	Ile	His	Leu	Gln	Glu	Leu	Asn	Leu	Asn	Asp	Asn	His
				125					130					135
Leu	Glu	Ser	Phe	Ser	Val	Ala	Leu	Cys	His	Ser	Thr	Leu	Gln	Lys
				140					145					150
Ser	Leu	Arg	Ser	Leu	Asp	Leu	Ser	Lys	Asn	Lys	Ile	Lys	Ala	Leu
				155					160					165
Pro	Val	Gln	Phe	Cys	Gln	Leu	Gln	Glu	Leu	Lys	Asn	Leu	Lys	Leu
				170					175					180
Asp	Asp	Asn	Glu	Leu	Ile	Gln	Phe	Pro	Cys	Lys	Ile	Gly	Gln	Leu
				185					190					195
Ile	Asn	Leu	Arg	Phe	Leu	Ser	Ala	Ala	Arg	Asn	Lys	Leu	Pro	Phe
				200					205					210
Leu	Pro	Ser	Glu	Phe	Arg	Asn	Leu	Ser	Leu	Glu	Tyr	Leu	Asp	Leu
				215					220					225
Phe	Gly	Asn	Thr	Phe	Glu	Gln	Pro	Lys	Val	Leu	Pro	Val	Ile	Lys
				230					235					240
Leu	Gln	Ala	Pro	Leu	Thr	Leu	Leu	Glu	Ser	Ser	Ala	Arg	Thr	Ile
				245					250					255
Leu	His	Asn	Arg	Ile	Pro	Tyr	Gly	Ser	His	Ile	Ile	Pro	Phe	His
				260					265					270
Leu	Cys	Gln	Asp	Leu	Asp	Thr	Ala	Lys	Ile	Cys	Val	Cys	Gly	Arg
				275					280					285
Phe	Cys	Leu	Asn	Ser	Phe	Ile	Gln	Gly	Thr	Thr	Thr	Met	Asn	Leu
				290					295					300
His	Ser	Val	Ala	His	Thr	Val	Val	Leu	Val	Asp	Asn	Leu	Gly	Gly
				305					310					315
Thr	Glu	Ala	Pro	Ile	Ile	Ser	Tyr	Phe	Cys	Ser	Leu	Gly	Cys	Tyr
				320					325					330
Val	Asn	Ser	Ser	Asp	Met	Leu	Lys							
				335										

<210> 10

<211> 164

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2960051CD1

<400> 10

Met	Lys	Ile	Ala	Val	Leu	Phe	Cys	Phe	Phe	Leu	Leu	Ile	Ile	Phe
1				5					10					15
Gln	Thr	Asp	Phe	Gly	Lys	Asn	Glu	Glu	Ile	Pro	Arg	Lys	Gln	Arg
				20					25					30
Arg	Lys	Ile	Tyr	His	Arg	Arg	Leu	Arg	Lys	Ser	Ser	Thr	Ser	His
				35					40					45
Lys	His	Arg	Ser	Asn	Arg	Gln	Leu	Gly	Ile	Pro	Gln	Thr	Thr	Val
				50					55					60
Phe	Thr	Pro	Val	Ala	Arg	Leu	Pro	Ile	Val	Asn	Phe	Asp	Tyr	Ser
				65					70					75
Met	Glu	Glu	Lys	Phe	Glu	Ser	Phe	Ser	Ser	Phe	Pro	Gly	Val	Glu
				80					85					90
Ser	Ser	Tyr	Asn	Val	Leu	Pro	Gly	Lys	Lys	Gly	His	Cys	Leu	Val
				95					100					105
Lys	Gly	Ile	Thr	Met	Tyr	Asn	Lys	Ala	Val	Trp	Ser	Pro	Glu	Pro
				110					115					120
Cys	Thr	Thr	Cys	Leu	Cys	Ser	Asp	Gly	Arg	Val	Leu	Cys	Asp	Glu
				125					130					135
Thr	Met	Cys	His	Pro	Gln	Arg	Cys	Pro	Gln	Thr	Val	Ile	Pro	Glu
				140					145					150
Gly	Glu	Cys	Cys	Pro	Val	Cys	Ser	Ala	Thr	Gly	Thr	Glu	Ile	
				155					160					

<210> 11

$\frac{1}{2} \left(\frac{1}{2} + \frac{1}{2} \right) = \frac{1}{2}$

WO 00/68380

PCT/US00/12811

Glu	Cys	Pro	Gln	Leu	Cys	Val	Cys	Glu	Ile	Arg	Pro	Trp	Phe	Thr
				35					40					45
Pro	Gln	Ser	Thr	Tyr	Arg	Glu	Ala	Thr	Thr	Val	Asp	Cys	Asn	Asp
				50					55					60
Leu	Arg	Leu	Thr	Arg	Ile	Pro	Ser	Asn	Leu	Ser	Ser	Asp	Thr	Gln
				65					70					75
Val	Leu	Leu	Leu	Gln	Ser	Asn	Asn	Ile	Ala	Lys	Thr	Val	Asp	Glu
				80					85					90
Leu	Gln	Gln	Leu	Phe	Asn	Leu	Thr	Glu	Leu	Asp	Phe	Ser	Gln	Asn
				95					100					105
Asn	Phe	Thr	Asn	Ile	Lys	Glu	Val	Gly	Leu	Ala	Asn	Leu	Thr	Gln
				110					115					120
Leu	Thr	Thr	Leu	His	Leu	Glu	Glu	Asn	Gln	Ile	Thr	Glu	Met	Thr
				125					130					135
Asp	Tyr	Cys	Leu	Gln	Asp	Leu	Ser	Asn	Leu	Gln	Glu	Leu	Tyr	Ile
				140					145					150
Asn	His	Asn	Gln	Ile	Ser	Thr	Ile	Ser	Ala	His	Ala	Phe	Ala	Gly
				155					160					165
Leu	Lys	Asn	Leu	Leu	Arg	Leu	His	Leu	Asn	Ser	Asn	Lys	Leu	Lys
				170					175					180
Val	Ile	Asp	Ser	Arg	Trp	Phe	Asp	Ser	Thr	Pro	Asn	Leu	Glu	Ile
				185					190					195
Leu	Met	Ile	Gly	Glu	Asn	Pro	Val	Ile	Gly	Ile	Leu	Asp	Met	Asn
				200					205					210
Phe	Lys	Pro	Leu	Ala	Asn	Leu	Arg	Ser	Leu	Val	Leu	Ala	Gly	Met
				215					220					225
Tyr	Leu	Thr	Asp	Ile	Pro	Gly	Asn	Ala	Leu	Val	Gly	Leu	Asp	Ser
				230					235					240
Leu	Glu	Ser	Leu	Ser	Phe	Tyr	Asp	Asn	Lys	Leu	Val	Lys	Val	Pro
				245					250					255
Gln	Leu	Ala	Leu	Gln	Lys	Val	Pro	Asn	Leu	Lys	Phe	Leu	Asp	Leu
				260					265					270
Asn	Lys	Asn	Pro	Ile	His	Lys	Ile	Gln	Glu	Gly	Asp	Phe	Lys	Asn
				275					280					285
Met	Leu	Arg	Leu	Lys	Glu	Leu	Gly	Ile	Asn	Asn	Met	Gly	Glu	Leu
				290					295					300
Val	Ser	Val	Asp	Arg	Tyr	Ala	Leu	Asp	Asn	Leu	Pro	Glu	Leu	Thr
				305					310					315
Lys	Leu	Glu	Ala	Thr	Asn	Asn	Pro	Lys	Leu	Ser	Tyr	Ile	His	Arg
				320					325					330
Leu	Ala	Phe	Arg	Ser	Val	Pro	Ala	Leu	Glu	Ser	Leu	Met	Leu	Asn
				335					340					345
Asn	Asn	Ala	Leu	Asn	Ala	Ile	Tyr	Gln	Lys	Thr	Val	Glu	Ser	Leu
				350					355					360
Pro	Asn	Leu	Arg	Glu	Ile	Ser	Ile	His	Ser	Asn	Pro	Leu	Arg	Cys
				365					370					375
Asp	Cys	Val	Ile	His	Trp	Ile	Asn	Ser	Asn</					

WO 00/68380

PCT/US00/12811

Val Ser Trp Lys	530	Val Asn Ser Asn Val	535	Thr Ser Asn Leu Lys	540
Trp Ser Ser Ala	545	Thr Met Lys Ile Asp	550	Asn Pro His Ile Thr Tyr	555
Thr Ala Arg Val	560	Pro Val Asp Val His	565	Glu Tyr Asn Leu Thr His	570
Leu Gln Pro Ser	575	Thr Asp Tyr Glu Val	580	Cys Leu Thr Val Ser Asn	585
Ile His Gln Gln	590	Thr Gln Lys Ser Cys	595	Val Asn Val Thr Thr Lys	600
Asn Ala Ala Phe	605	Ala Val Asp Ile Ser	610	Asp Gln Glu Thr Ser Thr	615
Ala Leu Ala Ala	620	Val Met Gly Ser Met	625	Phe Ala Val Ile Ser Leu	630
Ala Ser Ile Ala	635	Val Tyr Phe Ala Lys	640	Arg Phe Lys Arg Lys Asn	645
Tyr His His Ser	650	Leu Lys Lys Tyr Met	655	Gln Lys Thr Ser Ser Ile	660
Pro Leu Asn Glu	665	Leu Tyr Pro Pro Leu	670	Ile Asn Leu Trp Glu Gly	675
Asp Ser Glu Lys	680	Asp Lys Asp Gly Ser	685	Ala Asp Thr Lys Pro Thr	690
Gln Val Asp Thr	695	Ser Arg Ser Tyr Tyr	700	Met Trp	705
	710		715		

<210> 13

<211> 665

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4568384CD1

<400> 13

Met Val Leu Val Phe	5	His Lys Gly Glu Leu	10	Gly His Pro Leu Glu	15
Gln Ser Thr Asp Trp	20	Pro Lys Ser Pro Lys	25	Thr Pro Thr Gly Leu	30
Arg Arg Gly Arg Gln	35	Cys Ile Arg Pro Ala	40	Glu Ile Val Ala Ser	45
Leu Leu Glu Gly Glu	50	Glu Asn Thr Cys Gly	55	Lys Gln Lys Pro Lys	60
Glu Asn Asn Leu Lys	65	Pro Lys Phe Gln Ala	70	Phe Lys Gly Val Gly	75
Cys Leu Tyr Glu Lys	80	Glu Ser Met Lys Lys	85	Ser Leu Lys Asp Ser	90
Val Ala Ser Asn Asn	95	Lys Asp Gln Asn Ser	100	Met Lys His Glu Asp	105
Pro Ser Ile Ile Ser	110	Met Glu Asp Gly Ser	115	Pro Tyr Val Asn Gly	120
Ser Leu Gly Glu Val	125	Thr Pro Cys Gln His	130	Ala Lys Lys Ala Asn	135
Gly Pro Asn Tyr Ile	140	Gln Pro Gln Lys Arg	145	Gln Thr Thr Phe Glu	150
Ser Gln Asp Arg Lys	155	Ala Val Ser Pro Ser	160	Ser Ser Ser Glu Lys Arg	165
Ser Lys Asn Pro Ile	170	Ser Arg Pro Leu Glu	175	Gly Lys Lys Ser Leu	180
Ser Leu Ser Ala Lys	185	Thr His Asn Ile Gly	190	Phe Asp Lys Asp Ser	195
Cys His Ser Thr Thr	200	Lys Thr Glu Ala Ser	205	Gln Glu Glu Arg Ser	210
Asp Ser Ser Gly Leu	215	Thr Ser Leu Lys Lys	220	Ser Pro Lys Val Ser	225
Ser Lys Asp Thr Arg	230	Glu Ile Lys Thr Asp	235	Phe Ser Leu Ser Ile	240

WO 00/68380

PCT/US00/12811

Ser	Asn	Ser	Ser	Asp	Val	Ser	Ala	Lys	Asp	Lys	His	Ala	Glu	Asp
				245					250					255
Asn	Glu	Lys	Arg	Leu	Ala	Ala	Leu	Glu	Ala	Arg	Gln	Lys	Ala	Lys
				260					265					270
Glu	Val	Gln	Lys	Lys	Leu	Val	His	Asn	Ala	Leu	Ala	Asn	Leu	Asp
				275					280					285
Gly	His	Pro	Glu	Asp	Lys	Pro	Thr	His	Ile	Ile	Phe	Gly	Ser	Asp
				290					295					300
Ser	Glu	Cys	Glu	Thr	Glu	Glu	Thr	Ser	Thr	Gln	Glu	Gln	Ser	His
				305					310					315
Pro	Gly	Glu	Glu	Trp	Val	Lys	Glu	Ser	Met	Gly	Lys	Thr	Ser	Gly
				320					325					330
Lys	Leu	Phe	Asp	Ser	Ser	Asp	Asp	Asp	Glu	Ser	Asp	Ser	Glu	Asp
				335					340					345
Asp	Ser	Asn	Arg	Phe	Lys	Ile	Lys	Pro	Gln	Phe	Glu	Gly	Arg	Ala
				350					355					360
Gly	Gln	Lys	Leu	Met	Asp	Leu	Gln	Ser	His	Phe	Gly	Thr	Asp	Asp
				365					370					375
Arg	Phe	Arg	Met	Asp	Ser	Arg	Phe	Leu	Glu	Thr	Asp	Ser	Glu	Glu
				380					385					390
Glu	Gln	Glu	Glu	Val	Asn	Glu	Lys	Lys	Thr	Ala	Glu	Glu	Glu	Glu
				395					400					405
Leu	Ala	Glu	Glu	Lys	Lys	Lys	Ala	Leu	Asn	Val	Val	Gln	Ser	Val
				410					415					420
Leu	Gln	Ile	Asn	Leu	Ser	Asn	Ser	Thr	Asn	Arg	Gly	Ser	Val	Ala
				425					430					435
Ala	Lys	Lys	Phe	Lys	Asp	Ile	Ile	His	Tyr	Asp	Pro	Thr	Lys	Gln
				440					445					450
Asp	His	Ala	Thr	Tyr	Glu	Arg	Lys	Arg	Asp	Asp	Lys	Pro	Lys	Glu
				455					460					465
Ser	Lys	Ala	Lys	Arg	Lys	Lys	Lys	Arg	Glu	Glu	Ala	Glu	Lys	Leu
				470					475					480
Pro	Glu	Val	Ser	Lys	Glu	Met	Tyr	Tyr	Asn	Ile	Ala	Met	Asp	Leu
				485					490					495
Lys	Glu	Ile	Phe	Gln	Thr	Thr	Lys	Tyr	Thr	Ser	Glu	Lys	Glu	Glu
				500					505					510
Gly	Thr	Pro	Trp	Asn	Glu	Asp	Cys	Gly	Lys	Glu	Lys	Pro	Glu	Glu
				515					520					525
Ile	Gln	Asp	Pro	Ala	Ala	Leu	Thr	Ser	Asp	Ala	Glu	Gln	Pro	Ser
				530					535					540
Gly	Phe	Thr	Phe	Ser	Phe	Phe	Asp	Ser	Asp	Thr	Lys	Asp	Ile	Lys
				545					550					555
Glu	Glu	Thr	Tyr	Arg	Val	Glu	Thr	Val	Lys	Pro	Gly	Lys	Ile	Val
				560					565					570
Trp	Gln	Glu	Asp	Pro	Arg	Leu	Gln	Asp	Ser	Ser	Ser	Glu	Glu	Glu
				575					580					585
Asp	Val	Thr	Glu	Glu	Thr	Asp	His	Arg	Asn	Ser	Ser	Pro	Gly	Glu

```
<210> 14
<211> 547
<212> PRT
<213> Homo sapiens
```

```
<220>  
<221> misc_feature  
<223> Incyte ID No: 4586187CD1
```

[illegible]

WO 00/68380

PCT/US00/12811

<400>	14														
Met	Tyr	Ser	His	Asn	Val	Val	Ile	Met	Asn	Leu	Asn	Asn	Leu	Asn	
1				5					10						15
Leu	Thr	Gln	Val	Gln	Gln	Arg	Asn	Leu	Ile	Thr	Asn	Leu	Gln	Arg	
				20					25						30
Ser	Val	Asp	Asp	Thr	Ser	Gln	Ala	Ile	Gln	Arg	Ile	Lys	Asn	Asp	
				35					40						45
Phe	Gln	Asn	Leu	Gln	Gln	Val	Phe	Leu	Gln	Ala	Lys	Lys	Asp	Thr	
				50					55						60
Asp	Trp	Leu	Lys	Glu	Lys	Val	Gln	Ser	Leu	Gln	Thr	Leu	Ala	Ala	
				65					70						75
Asn	Asn	Ser	Ala	Leu	Ala	Lys	Ala	Asn	Asn	Asp	Thr	Leu	Glu	Asp	
				80					85						90
Met	Asn	Ser	Gln	Leu	Asn	Ser	Phe	Thr	Gly	Gln	Met	Glu	Asn	Ile	
				95					100						105
Thr	Thr	Ile	Ser	Gln	Ala	Asn	Glu	Gln	Asn	Leu	Lys	Asp	Leu	Gln	
				110					115						120
Asp	Leu	His	Lys	Asp	Ala	Glu	Asn	Arg	Thr	Ala	Ile	Lys	Phe	Asn	
				125					130						135
Gln	Leu	Glu	Glu	Arg	Phe	Gln	Leu	Phe	Glu	Thr	Asp	Ile	Val	Asn	
				140					145						150
Ile	Ile	Ser	Asn	Ile	Ser	Tyr	Thr	Ala	His	His	Leu	Arg	Thr	Leu	
				155					160						165
Thr	Ser	Asn	Leu	Asn	Glu	Val	Arg	Thr	Thr	Cys	Thr	Asp	Thr	Leu	
				170					175						180
Thr	Lys	His	Thr	Asp	Asp	Leu	Thr	Ser	Leu	Asn	Asn	Thr	Leu	Ala	
				185					190						195
Asn	Ile	Arg	Leu	Asp	Ser	Val	Ser	Leu	Arg	Met	Gln	Gln	Asp	Leu	
				200					205						210
Met	Arg	Ser	Arg	Leu	Asp	Thr	Glu	Val	Ala	Asn	Leu	Ser	Val	Ile	
				215					220						225
Met	Glu	Glu	Met	Lys	Leu	Val	Asp	Ser	Lys	His	Gly	Gln	Leu	Ile	
				230					235						240
Lys	Asn	Phe	Thr	Ile	Leu	Gln	Gly	Pro	Pro	Gly	Pro	Arg	Gly	Pro	
				245					250						255
Arg	Gly	Asp	Arg	Gly	Ser	Gln	Gly	Pro	Pro	Gly	Pro	Thr	Gly	Asn	
				260					265						270
Lys	Gly	Gln	Lys	Gly	Glu	Lys	Gly	Glu	Pro	Gly	Pro	Pro	Gly	Pro	
				275					280						285
Ala	Gly	Glu	Arg	Gly	Pro	Ile	Gly	Pro	Ala	Gly	Pro	Pro	Gly	Glu	
				290					295						300
Arg	Gly	Gly	Lys	Gly	Ser	Lys	Gly	Ser	Gln	Gly	Pro	Lys	Gly	Ser	
				305					310						315
Arg	Gly	Ser	Pro	Gly	Lys	Pro	Gly	Pro	Gln	Gly	Pro	Ser	Gly	Asp	
				320					325						330
Pro	Gly	Pro													

WO 00/68380

PCT/US00/12811

Lys Ala Gly Gln Pro Asp Asn Trp Gly His Gly His Gly Pro Gly
 500 505 510
 Glu Asp Cys Ala Gly Leu Ile Tyr Ala Gly Gln Trp Asn Asp Phe
 515 520 525
 Gln Cys Glu Asp Val Asn Asn Phe Ile Cys Glu Lys Asp Arg Glu
 530 535 540
 Thr Val Leu Ser Ser Ala Leu
 545

<210> 15
 <211> 109
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 401801CD1

<400> 15
 Met Tyr Phe Asn Leu Gln Glu Asn Ile Phe Met Tyr Gly Gly Arg
 1 5 10 15
 Ile Glu Thr Asn Asp Gly Asn Val Thr Asp Glu Leu Trp Val Phe
 20 25 30
 Asn Ile His Ser Gln Ser Trp Ser Thr Lys Thr Pro Thr Val Leu
 35 40 45
 Gly His Gly Gln Gln Tyr Ala Val Glu Gly His Ser Ala His Ile
 50 55 60
 Met Glu Leu Asp Ser Arg Asp Val Val Met Ile Ile Ile Phe Gly
 65 70 75
 Tyr Ser Ala Ile Tyr Gly Tyr Thr Ser Ser Ile Gln Glu Tyr His
 80 85 90
 Ile Cys Glu Leu Leu Lys Asn Cys Asn Phe Phe Ile Asp Trp Glu
 95 100 105
 Cys Phe Ser Leu

<210> 16
 <211> 192
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1721842CD1

<400> 16
 Met Asn Lys Arg Asp Tyr Met Asn Thr Ser Val Gln Glu Pro Pro
 1 5 10 15
 Leu Asp Tyr Ser Phe Arg Ser Ile His Val Ile Gln Asp Leu Val
 20 25 30
 Asn Glu Glu Pro Arg Thr Gly Leu Arg Pro Leu Lys Arg Ser Lys
 35 40 45
 Ser Gly Lys Ser Leu Thr Gln Ser Leu Trp Leu Asn Asn Asn Val
 50 55 60
 Leu Asn Asp Leu Arg Asp Phe Asn Gln Val Ala Ser Gln Leu Leu
 65 70 75
 Glu His Pro Glu Asn Leu Ala Trp Ile Asp Leu Ser Phe Asn Asp
 80 85 90
 Leu Thr Ser Ile Asp Pro Val Leu Thr Thr Phe Phe Asn Leu Ser
 95 100 105
 Val Leu Tyr Leu His Gly Asn Ser Ile Gln Arg Leu Gly Glu Val
 110 115 120
 Asn Lys Leu Ala Val Leu Pro Arg Leu Arg Ser Leu Thr Leu His
 125 130 135
 Gly Asn Pro Met Glu Glu Lys Gly Tyr Arg Gln Tyr Val Leu
 140 145 150
 Cys Thr Leu Ser Arg Ile Thr Thr Phe Asp Phe Ser Gly Val Thr
 155 160 165
 Lys Ala Asp Arg Thr Thr Ala Glu Val Trp Lys Arg Met Asn Ile

WO 00/68380

PCT/US00/12811

170
Lys Pro Lys Lys Ala Trp Thr Lys Gln Asn Thr Leu
185 190

180

<210> 17
<211> 575
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1833221CD1

<400> 17
Met Val Leu Gly Ser Phe Gly Thr Asp Leu Met Arg Glu Arg Arg
1 5 10 15
Asp Leu Glu Arg Arg Thr Asp Ser Ser Ile Ser Asn Leu Met Asp
20 25 30
Tyr Ser His Arg Ser Gly Asp Phe Thr Thr Ser Ser Tyr Val Gln
35 40 45
Asp Arg Val Pro Ser Tyr Ser Gln Gly Ala Arg Pro Lys Glu Asn
50 55 60
Ser Met Ser Thr Leu Gln Leu Asn Thr Ser Ser Thr Asn His Gln
65 70 75
Leu Pro Ser Glu His Gln Thr Ile Leu Ser Ser Arg Asp Ser Arg
80 85 90
Asn Ser Leu Arg Ser Asn Phe Ser Ser Arg Glu Ser Glu Ser Ser
95 100 105
Arg Ser Asn Thr Gln Pro Gly Phe Ser Tyr Ser Ser Ser Arg Asp
110 115 120
Glu Ala Pro Ile Ile Ser Asn Ser Glu Arg Val Val Ser Ser Gln
125 130 135
Arg Pro Phe Gln Glu Ser Ser Asp Asn Glu Gly Arg Arg Thr Thr
140 145 150
Arg Arg Leu Leu Ser Arg Ile Ala Ser Ser Met Ser Ser Thr Phe
155 160 165
Phe Ser Arg Arg Ser Ser Gln Asp Ser Leu Asn Thr Arg Ser Leu
170 175 180
Asn Ser Glu Asn Ser Tyr Val Ser Pro Arg Ile Leu Thr Ala Ser
185 190 195
Gln Ser Arg Ser Asn Val Pro Ser Ala Ser Glu Val Pro Asp Asn
200 205 210
Arg Ala Ser Glu Ala Ser Gln Gly Phe Arg Phe Leu Arg Arg Arg
215 220 225
Trp Gly Leu Ser Ser Leu Ser His Asn His Ser Ser Glu Ser Asp
230 235 240
Ser Glu Asn Phe Asn Gln Glu Ser Glu Gly Arg Asn Thr Gly Pro
245 250 255
Trp Leu Ser Ser Ser Leu Arg Asn Arg Cys Thr Pro Leu Phe Ser
260 265 270
Arg Arg Arg Arg Glu Gly Arg Asp Glu Ser Ser Arg Ile Pro Thr
275 280 285
Ser Asp Thr Ser Ser Arg Ser His Ile Phe Arg Arg Glu Ser Asn
290 295 300
Glu Val Val His Leu Glu Ala Gln Asn Asp Pro Leu Gly Ala Ala
305 310 315
Ala Asn Arg Pro Gln Ala Ser Ala Ala Ser Ser Ser Ala Thr Thr
320 325 330
Gly Gly Ser Thr Ser Asp Ser Ala Gln Gly Gly Arg Asn Thr Gly
335 340 345
Ile Ser Gly Ile Leu Pro Gly Ser Leu Phe Arg Phe Ala Val Pro
350 355 360
Pro Ala Leu Gly Ser Asn Leu Thr Asp Asn Val Met Ile Thr Val
365 370 375
Asp Ile Ile Pro Ser Gly Trp Asn Ser Ala Asp Gly Lys Ser Asp
380 385 390
Lys Thr Lys Ser Ala Pro Ser Arg Asp Pro Glu Arg Leu Gln Lys
395 400 405

WO 00/68380

PCT/US00/12811

Ile	Lys	Glu	Ser	Leu	Leu	Glu	Asp	Ser	Glu	Glu	Glu	Glu	Gly	
				410				415					420	
Asp	Leu	Cys	Arg	Ile	Cys	Gln	Met	Ala	Ala	Ser	Ser	Ser	Asn	
				425				430					435	
Leu	Leu	Ile	Glu	Pro	Cys	Lys	Cys	Thr	Gly	Ser	Leu	Gln	Tyr	Val
				440				445					450	
His	Gln	Asp	Cys	Met	Lys	Lys	Trp	Leu	Gln	Ala	Lys	Ile	Asn	Ser
				455				460					465	
Gly	Ser	Ser	Leu	Glu	Ala	Val	Thr	Thr	Cys	Glu	Leu	Cys	Lys	Glu
				470				475					480	
Lys	Leu	Glu	Leu	Asn	Leu	Glu	Asp	Phe	Asp	Ile	His	Glu	Leu	His
				485				490					495	
Arg	Ala	His	Ala	Asn	Glu	Gln	Ala	Glu	Tyr	Glu	Phe	Ile	Ser	Ser
				500				505					510	
Gly	Leu	Tyr	Leu	Val	Val	Leu	Leu	His	Leu	Cys	Glu	Gln	Ser	Phe
				515				520					525	
Ser	Asp	Met	Met	Gly	Asn	Thr	Asn	Glu	Pro	Ser	Thr	Arg	Val	Arg
				530				535					540	
Phe	Ile	Asn	Leu	Ala	Arg	Thr	Leu	Gln	Ala	His	Met	Glu	Asp	Leu
				545				550					555	
Glu	Thr	Ser	Glu	Asp	Asp	Ser	Glu	Glu	Asp	Gly	Asp	His	Asn	Arg
				560				565					570	
Thr	Phe	Asp	Ile	Ala										
				575										

<210> 18

<211> 342

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2041168CD1

<400> 18

Met	Ala	Glu	Gly	Gly	Ser	Gly	Asp	Val	Asp	Asp	Ala	Gly	Asp	Cys	
1				5					10					15	
Ser	Gly	Ala	Arg	Tyr	Asn	Asp	Trp	Ser	Asp	Asp	Asp	Asp	Asp	Ser	
				20					25					30	
Asn	Glu	Ser	Lys	Ser	Ile	Val	Trp	Tyr	Pro	Pro	Trp	Ala	Arg	Ile	
				35					40					45	
Gly	Thr	Glu	Ala	Gly	Thr	Arg	Ala	Arg	Ala	Arg	Ala	Arg	Ala	Arg	
				50					55					60	
Ala	Thr	Arg	Ala	Arg	Arg	Ala	Val	Gln	Lys	Arg	Ala	Ser	Pro	Asn	
				65					70					75	
Ser	Asp	Asp	Thr	Val	Leu	Ser	Pro	Gln	Glu	Leu	Gln	Lys	Val	Leu	
				80					85					90	
Cys	Leu	Val	Glu	Met	Ser	Glu	Lys	Pro	Tyr	Ile	Leu	Glu	Ala	Ala	
				95					100					105	
Leu	Ile	Ala	Leu	Gly	Asn	Asn	Ala	Ala	Tyr	Ala	Phe	Asn	Arg	Asp	
				110					115					120	
Ile	Ile	Arg	Asp	Leu	Gly	Gly	Leu	Pro	Ile	Val	Ala	Lys	Ile	Leu	
				125					130					135	
Asn	Thr	Arg	Asp	Pro	Ile	Val	Lys	Glu	Lys	Ala	Leu	Ile	Val	Leu	
				140					145					150	
Asn	Asn	Leu	Ser	Val	Asn	Ala	Glu	Asn	Gln	Arg	Arg	Leu	Lys	Val	
				155					160					165	
Tyr	Met	Asn	Gln	Val	Cys	Asp	Asp	Thr	Ile	Thr	Ser	Arg	Leu	Asn	
				170					175					180	
Ser	Ser	Val	Gln	Leu	Ala	Gly	Leu	Arg	Leu	Leu	Thr	Asn	Met	Thr	
				185					190					195	
Val	Thr	Asn	Glu	Tyr	Gln	His	Met	Leu	Ala	Asn	Ser	Ile	Ser	Asp	
				200					205					210	
Phe	Phe	Arg	Leu	Phe	Ser	Ala	Gly	Asn	Glu	Glu	Thr	Lys	Leu	Gln	
				215					220					225	
Val	Leu	Lys	Leu	Leu	Leu	Asn	Leu	Ala	Glu	Asn	Pro	Ala	Met	Thr	
				230					235					240	
Arg	Glu	Leu	Leu	Arg	Ala	Gln	Val	Pro	Ser	Ser	Leu	Gly	Ser	Leu	

WO 00/68380

PCT/US00/12811

Gly Leu Glu Asp Glu Ser Gly Leu Val Glu Leu Glu Leu Arg Gly
 155 160 165
 Glu Met Leu Thr Gly Thr Gly
 170

<210> 23

<211> 571

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2850987CD1

<400> 23

Met Thr Arg Ala Gly Asp His Asn Arg Gln Arg Gly Cys Cys Gly
 1 5 10 15
 Ser Leu Ala Asp Tyr Leu Thr Ser Ala Lys Phe Leu Leu Tyr Leu
 20 25 30
 Gly His Ser Leu Ser Thr Trp Gly Asp Arg Met Trp His Phe Ala
 35 40 45
 Val Ser Val Phe Leu Val Glu Leu Tyr Gly Asn Ser Leu Leu Leu
 50 55 60
 Thr Ala Val Tyr Gly Leu Val Val Ala Gly Ser Val Leu Val Leu
 65 70 75
 Gly Ala Ile Ile Gly Asp Trp Val Asp Lys Asn Ala Arg Leu Lys
 80 85 90
 Val Ala Gln Thr Ser Leu Val Val Gln Asn Val Ser Val Ile Leu
 95 100 105
 Cys Gly Ile Ile Leu Met Met Val Phe Leu His Lys His Glu Leu
 110 115 120
 Leu Thr Met Tyr His Gly Trp Val Leu Thr Ser Cys Tyr Ile Leu
 125 130 135
 Ile Ile Thr Ile Ala Asn Ile Ala Asn Leu Ala Ser Thr Ala Thr
 140 145 150
 Ala Ile Thr Ile Gln Arg Asp Trp Ile Val Val Val Ala Gly Glu
 155 160 165
 Asp Arg Ser Lys Leu Ala Asn Met Asn Ala Thr Ile Arg Arg Ile
 170 175 180
 Asp Gln Leu Thr Asn Ile Leu Ala Pro Met Ala Val Gly Gln Ile
 185 190 195
 Met Thr Phe Gly Ser Pro Val Ile Gly Cys Gly Phe Ile Ser Gly
 200 205 210
 Trp Asn Leu Val Ser Met Cys Val Glu Tyr Val Leu Leu Trp Lys
 215 220 225
 Val Tyr Gln Lys Thr Pro Ala Leu Ala Val Lys Ala Gly Leu Lys
 230 235 240
 Glu Glu Glu Thr Glu Leu Lys Gln Leu Asn Leu His Lys Asp Thr
 245 250 255
 Glu Pro Lys Pro Leu Glu Gly Thr His Leu Met Gly Val Lys Asp
 260 265 270
 Ser Asn Ile His Glu Leu Glu His Glu Gln Glu Pro Thr Cys Ala
 275 280 285
 Ser Gln Met Ala Glu Pro Phe Arg Thr Phe Arg Asp Gly Trp Val
 290 295 300
 Ser Tyr Tyr Asn Gln Pro Val Phe Leu Ala Gly Met Gly Leu Ala
 305 310 315
 Phe Leu Tyr Met Thr Val Leu Gly Phe Asp Cys Ile Thr Thr Gly
 320 325 330
 Tyr Ala Tyr Thr Gln Gly Leu Ser Gly Ser Ile Leu Ser Ile Leu
 335 340 345
 Met Gly Ala Ser Ala Ile Thr Gly Ile Met Gly Thr Val Ala Phe
 350 355 360
 Thr Trp Leu Arg Lys Cys Gly Leu Val Arg Thr Gly Leu Ile
 365 370 375
 Ser Gly Leu Ala Gln Leu Ser Cys Leu Ile Leu Cys Val Ile Ser
 380 385 390
 Val Phe Met Pro Gly Ser Pro Leu Asp Leu Ser Val Ser Pro Phe

WO 00/68380

PCT/US00/12811

	395		400		405
Glu Asp Ile Arg	Ser Arg Phe Ile Gln	Gly Glu Ser Ile Thr	Pro		
	410		415		420
Thr Lys Ile Pro	Glu Ile Thr Thr Glu	Ile Tyr Met Ser Asn	Gly		
	425		430		435
Ser Asn Ser Ala	Asn Ile Val Pro Glu	Thr Ser Pro Glu Ser	Val		
	440		445		450
Pro Ile Ile Ser	Val Ser Leu Leu Phe	Ala Gly Val Ile Ala	Ala		
	455		460		465
Arg Ile Gly Leu	Trp Ser Phe Asp Leu	Thr Val Thr Gln Leu	Leu		
	470		475		480
Gln Glu Asn Val	Ile Glu Ser Glu Arg	Gly Ile Ile Asn Gly	Val		
	485		490		495
Gln Asn Ser Met	Asn Tyr Leu Leu Asp	Leu Leu His Phe Ile	Met		
	500		505		510
Val Ile Leu Ala	Pro Asn Pro Glu Ala	Phe Gly Leu Leu Val	Leu		
	515		520		525
Ile Ser Val Ser	Phe Val Ala Met Gly	His Ile Met Tyr Phe	Arg		
	530		535		540
Phe Ala Gln Asn	Thr Leu Gly Asn Lys	Leu Phe Ala Cys Gly	Pro		
	545		550		555
Asp Ala Lys Glu	Val Arg Lys Glu Asn	Gln Ala Asn Thr Ser	Val		
	560		565		570

Val

<210> 24

<211> 455

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3557211CD1

<400> 24

Met Asp Pro Thr	Gly Asn Ser Ala Thr	Pro Gln Ile Leu Glu Leu	
1	5	10	15
Lys Trp Ser His	Ile Glu Trp Ser Gln Thr	Glu Tyr Ile Cys Glu	
	20	25	30
Asn Val Gly Leu	Leu Pro Leu Glu Ile Ile	Arg Arg Gly Tyr Ser	
	35	40	45
Met Asp Ser Ala	Phe Val Gly Ile Lys Val	Asn Gln Val Ser Ala	
	50	55	60
Ala Val Gly Lys	Asp Phe Thr Val Ile Pro	Ser Lys Leu Ile Gln	
	65	70	75
Phe Asp Pro Gly	Met Ser Thr Lys Met Trp	Asn Ile Ala Ile Thr	
	80	85	90
Tyr Asp Gly Leu	Glu Glu Asp Asp Glu Val	Phe Glu Val Ile Leu	
	95	100	105
Asn Ser Pro Val	Asn Ala Val Leu Gly Thr	Lys Thr Lys Ala Ala	
	110	115	120
Val Lys Ile Leu	Asp Ser Lys Gly Gly Gln	Cys His Pro Ser Tyr	
	125	130	135
Ser Ser Asn Gln	Ser Lys His Ser Thr Trp	Glu Lys Gly Ile Trp	
	140	145	150
His Leu Leu Pro	Pro Gly Ser Ser Ser Ser	Thr Thr Ser Gly Ser	
	155	160	165
Phe His Leu Glu	Arg Arg Pro Leu Pro Ser	Ser Met Gln Leu Ala	
	170	175	180
Val Ile Arg Gly	Asp Thr Leu Arg Gly Phe	Asp Ser Thr Asp Leu	
	185	190	195
Ser Gln Arg Lys	Leu Arg Thr Arg Gly Asn	Gly Lys Thr Val Arg	
	200	205	210
Pro Ser Ser Val	Tyr Arg Asn Gly Thr Asp	Ile Ile Tyr Asn Tyr	
	215	220	225
His Gly Ile Val	Ser Leu Lys Leu Glu Asp	Asp Ser Phe Pro Thr	
	230	235	240
His Lys Arg Lys	Ala Lys Val Ser Ile Ile	Ser Gln Pro Gln Lys	

WO 00/68380

PCT/US00/12811

Ser	Asn	Trp	Glu	Gln	Ile	Leu	Tyr	Val	Thr	Glu	Pro	Glu	Ala	Trp
				215					220					225
Thr	Ala	Ala	Ala	Met	Tyr	Gln	Ala	Thr	Arg	Ile	Phe	Ala	Ser	Asn
				230					235					240
Leu	Lys	Glu	Arg	Met	Ala	Gln	Arg	Phe	Tyr	Asn	Leu	Val	Leu	Leu
				245					250					255
Pro	Arg	Val	Arg	Asp	Asp	Val	Ala	Glu	Tyr	Lys	Arg	Leu	Asn	Phe
				260					265					270
His	Leu	Tyr	Met	Ala	Leu	Lys	Lys	Ala	Leu	Phe	Lys	Pro	Gly	Ala
				275					280					285
Trp	Phe	Lys	Gly	Ile	Leu	Ile	Pro	Leu	Cys	Glu	Ser	Gly	Thr	Cys
				290					295					300
Thr	Leu	Arg	Glu	Ala	Ile	Ile	Val	Gly	Ser	Ile	Ile	Thr	Lys	Cys
				305					310					315
Ser	Ile	Pro	Val	Leu	His	Ser	Ser	Ala	Ala	Met	Leu	Lys	Ile	Ala
				320					325					330
Glu	Met	Glu	Tyr	Ser	Gly	Ala	Asn	Ser	Ile	Phe	Leu	Arg	Leu	Leu
				335					340					345
Leu	Asp	Lys	Lys	Tyr	Ala	Leu	Pro	Tyr	Arg	Val	Leu	Asp	Ala	Leu
				350					355					360
Val	Phe	His	Phe	Leu	Gly	Phe	Arg	Thr	Glu	Lys	Arg	Glu	Leu	Pro
				365					370					375
Val	Leu	Trp	His	Gln	Cys	Leu	Leu	Thr	Leu	Val	Gln	Arg	Tyr	Lys
				380					385					390
Ala	Asp	Leu	Ala	Thr	Asp	Gln	Lys	Glu	Ala	Leu	Leu	Glu	Leu	Leu
				395					400					405
Arg	Leu	Gln	Pro	His	Pro	Gln	Leu	Ser	Pro	Glu	Ile	Arg	Arg	Glu
				410					415					420
Leu	Gln	Ser	Ala	Val	Pro	Arg	Asp	Val	Glu	Asp	Val	Pro	Ile	Thr
				425					430					435

Val Glu

<210> 26

<211> 2893

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 398269CB1

<400> 26

agtggctgag	tcgggggcgg	gctgggaggg	ctgtcggtgg	gccagtcctgc	gtacgacggc	60
ccgtcccctg	cgcacggacg	ccgggaagaa	gggggtgggg	ccacgtttgc	gtccgcgcca	120
tcaggcccg	gatagcggcg	agggtccgctt	tcagtgtatg	gttttccctg	ccaaacgggt	180
ctgcttggtg	ccatccatgg	agggcggtgcg	ctgggccttt	tcctgcggca	cttggctgcc	240
gagccgagcc	gaatggctgc	tggcagtgcg	atcgattcag	cccgaggaga	aggagcgc	300
tggccagttc	gtctttgccc	gggacgctaa	ggcagccatg	gctggtcgtc	tgatgataag	360
gaaattagtt	gcagagaaat	tgaatatccc	ttggaatcat	attcgtttgc	aaagaactgc	420
aaaaggaaaa	ccagttcttg	caaaggactc	atcgaatcct	taccgaatt	tcaactttaa	480
catctctcat	caaggagact	atgcagtgct	tgtctgtgaa	cctgagctgc	aagttggaat	540
tgatataatg	aagactagtt	ttccaggctg	tggttcaatt	ccagaattct	ttcatattat	600
gaaaagaaaag	tttaccaaca	aagaatggga	aacaatcaga	agctttaagg	atgagtgga	660
tcagctggat	atgttttata	ggaattgggc	acttaaggaa	agcttcataa	aagccattgg	720
tgttggacta	ggatttgaat	tgcagcggct	tgaatttgat	ctatctccat	taaacttgga	780
tataggccaa	gtttataaag	aaacacgttt	attcctggat	ggagaggaag	aaaaagaatg	840
ggcatttgag	gaaagcaaaa	tagatgagca	ccattttggt	gcagttgctc	ttaggaaacc	900
cgatggatct	agacatcagg	atgttccatc	tcaggatgat	tccaaaccaa	cccagaggca	960
atttactatt	ctcaacttta	atgatttaat	gtcatctgcc	gttcccatga	cacctgaaga	1020
tccttcattt	tgggactggt	tttgcttcac	agaagaaatt	ccaatacgaa	atggtacaaa	1080
gtcatgatga	ttccctgagt	aacaaaggga	aatgaaaact	gtttgtgatc	ttccgtattc	1140
actgaaaaat	aaatgcttgt	ttagtatcaa	attttatttc	acgaaagttt	ttttaaagaa	1200
cagaaaacttt	tccaattaaa	aaaaaaaaagc	agactctctg	ttcaagatag	ctcactggaa	1260
tacatgttta	cctctttctt	tcctaaattg	cattgaattg	ataggaagga	tggcgggaatc	1320
ttaaagtgat	acatgctaac	tgtagaaaaa	aatagaaaat	gcacataagc	aaaaggaaac	1380
attttaaagtc	tatctttcaa	agataactac	tcttaaaacc	ttgagtatct	tttcagacct	1440
ttttcttggc	aaatgaatcc	atattgacat	atttgatttt	tttaaaaaca	tggaaacgta	1500

WO 00/68380

PCT/US00/12811

ctgtttttgta	aattctttttt	aactgcacat	ctactgttca	taaatatacc	tctgtaacat	1560
aacttttttgt	ggttctaata	tactctgttg	tatagctaat	acaagttagg	atgctttttgg	1620
ccagaggtaa	cagtgctcaa	atataattgg	cctaagtaac	ctaggaaatt	gtttgcacata	1680
acacagggtt	cagggtgtgc	attaaagaca	cactttttttt	gccttgacct	cagttgggttt	1740
gttttgccct	aggctattcc	acctctcgat	cacaagagct	gctgctataa	cttccagttt	1800
tacatctttg	taaaattgta	tctcaaaggc	agaaaaggcc	atttcgtctc	tcattttgttt	1860
tatccatgag	gaagattttt	acaaaaagcc	tccagaagat	ttccccctcg	tttccattga	1920
cttagatcag	gttacagaga	aaggcaatgt	ctgacatttt	tggctctctg	tagaagtaga	1980
ctctgttgaa	aagaaagaag	ctaagctagg	tgtgaagaat	ggaattggaa	gccactgccc	2040
ttcccataag	aaaggttttac	cataattttac	tcactttttt	ctgtgttata	catttttgatt	2100
atctgcagtt	tattactaca	agcagtggca	gagtgaattgt	ccttgtacat	tttgagttac	2160
atgtcttaatt	atgtccttga	gaaagttttct	aaaagtggaa	tgattggttt	gactggttca	2220
tagggccttta	attatacaat	ttacccctct	aattagtact	atatgtatgt	gacttccctc	2280
cccttgcag	aatactcctt	ggtcaattgt	aggttattct	tttggtttaa	tttttgccaa	2340
tgtaattaaa	aaatggtatg	tcatttttta	aatttgtatt	tctttcatta	caataaagat	2400
tgttatgtca	gtattgttat	tggccttttcg	tattctctct	aacgtgaacc	gtctgttcac	2460
tgtttttacc	tgttttcgtt	ttagcaagta	gtacttaatt	taaagtgtga	acttaatat	2520
taagatgcc	ggaccatcat	attgatgaca	aaaatctatt	atgtgggttt	agtgcatgtc	2580
tggagtttaa	actcttaagt	atcaagaagt	aaaataattt	cattgtttct	gttactaaca	2640
ttaaaagtgt	gcaaattgta	taaaattagg	tacttcgatt	agtaaaagaa	aaaaatttga	2700
atgttttttt	ttttatttta	ttaagcatgc	ccagttatgc	caagcatagt	aaataaagggt	2760
caagtagcat	ttataataga	ggaagtattg	ttatccctag	catgagtgt	atggtgat	2820
gaaaaacttt	gtctgtcat	tataataata	aaaaaatgaa	catttattat	ggaatttcaa	2880
aaaaaaaaaa	aaa					2893

```
<210> 27
<211> 2276
<212> DNA
<213> Homo sapiens
```

```
<220>  
<221> misc_feature  
<223> Incyte ID No: 1258888CB1
```

<400>	27					
gcgagtgagg	cggaggaccc	gagcggtga	ggagagagga	ggcggcggt	tagctgctac	60
ggggtcgggc	cggcgccctc	ccgagggggg	ctcaggagga	ggaaaggagga	cccgtgcgag	120
aatgcctctc	ccctggagcc	ttgcgtctcc	ctgctgtctc	tcctgggtgg	caggtgggtt	180
cggaacgcg	gccagtcaa	ggcatcacgg	gttgttagca	tcggcacgtc	agcctggggt	240
ctgtcactat	ggaactaaac	tggcctgtcg	ctacggctgg	agaagaaaca	gcaagggagt	300
ctgtgaagct	acatgcgaac	ctggatgtaa	gtttgggtgag	tgcgtgggac	caaacaaatg	360
catgtgcctt	ccaggatata	ccgggaaaac	ctgcagtcac	gatgtgaatg	agtgtggaat	420
gaaaccccg	ccatgccaac	acagatgtgt	gaatacacac	ggaaagctaca	agtgcttttg	480
cctcagtggc	cacatgtctc	tgccagatgc	tacgtgtgtg	aactctagga	catgtgccat	540
gataaaactgt	cagtcagctc	ctaagacac	agaagaaggg	ccacagtggc	tgtgtccatc	600
ctcaggactc	cgctggccc	caaatggaag	agactgtcta	gatattgatg	aatgtgcctc	660
tggtaaagtc	atctgtccct	acaatcgaag	atgtgtgaac	acatttgga	gctactactg	720
caaatgtcac	attgtttcgc	aactgcaata	tatcagtgga	cgatatgact	gtatagatat	780
aaatgaatgt	actatggata	gccatacgtg	cagccaccat	gccaattgct	tcaataccca	840
aggttccttc	aagtgtaaat	gcaagcaggg	atataaaggc	aatggacttc	gggtttctgc	900
tatccctgaa	aattctgtga	aggaagtctc	cagagcacct	ggtaccatca	aagacagaat	960
caagaagttg	cttgctcaca	aaaacagcat	gaaaaagaag	gcaaaaatta	aaaatgttac	1020
cccgaaccc	accaggactc	ctacccttaa	ggtgaacttg	cagcccttca	actatgaaga	1080
gatagtttcc	agaggcggga	actctcatgg	aggtaaaaaa	gggaatgaag	agaaaaatga	1140
agaggggctt	gaggatgaga	aaagagaaga	gaaagccctg	agaatgaca	tagaggagcg	1200
aagcctgcga	ggagatgtgt	ttttccctaa	ggtgaatgaa	gcaggtgaat	tcggcctgat	1260
tctggctcaa	aggaaaagcgc	taacttccaa	actggaacat	aaagcagatg	taaatacttc	1320
ggttgactgc	agcttcaatc	atgggatctg	tgactggaaa	caggatagag	aagatgattt	1380
tgactggaat	cctgctgata	gagataatgc	tattggcttc	tatatggcag	ttccggcctt	1440
ggcaggtcac	aagaaagaca	ttggccgatt	gaaacttctc	ctacctgacc	tgcaacccca	1500
aagcaacttc	tgtttgtctt	ttgattaccg	gctggccgga	gacaaagtgc	ggaaacttcg	1560
agtgtttgtg	aaaaacagta	acaattgccct	ggcatgggag	agagccacga	gtgaggatga	1620
aaagttggaag	acagggaaaa	ttcagttgta	tcaaggaaact	gatgtacca	aaagcatcat	1680
ttttgaagca	gaacgtggca	agggcaaaac	cggcgaaatc	gcagtggatg	gcgtcttgct	1740
tgtttcaggc	ttatgtccag	atagcctttt	atctgtggat	gactgaatgt	tactactctt	1800
atatttgact	ttgtattgtc	gttccctggg	ttttttgata	tgtcatcata	ggacctctgg	1860
cattttagaa	ttactagctg	aaaaattgta	atgtaccaac	agaaatatta	ttgtaaagatg	1920

WO 00/68380

PCT/US00/12811

cctttcttgt	ataagatatg	ccaatatattg	ccttaaatat	catatcactg	tatcttctca	1980
gtcattttctg	aattctttcca	cattatatta	taaaatatgg	aaatgtcagt	ttatctcccc	2040
tcctcagtat	atctgatttg	tataagtaag	ttgatgagct	tctctctaca	acatttctag	2100
aaaatagaaa	aaaaagcaca	gagaaatgtt	taactgtttg	actcttatga	tacttcttgg	2160
aaactatgac	atcaaagata	gactttttgcc	taagtggctt	agctgggtct	ttcatagcca	2220
aacttgtata	tttaaattct	ttgtaataat	aatatccaaa	tcataaaaaa	aaaaaa	2276

<210> 28
 <211> 2016
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1375891CB1

<400> 28						
gaaaagggtac	ccgcgagaga	cagccagcag	ttctgtggag	cagcgggtggc	cggctaggat	60
gggctgtctc	tgggggtctg	ctctgcccc	tttcttcttc	tgctgggagg	ttgggggtctc	120
tgggagctct	gcaggcccc	gcaccccgag	agcagacact	gcgatgacaa	cggacgacac	180
agaagtgtcc	gctatgactc	tagcaccggg	ccacgccgt	ctggaaaactc	aaacgctgag	240
cgtgagacc	tcttctaggg	cctcaacccc	agccggcccc	attccagaag	cagagaccag	300
gggagccaag	agaatttccc	ctgcaagaga	gaccaggagt	ttcacaaaaa	catctcccaa	360
cttcatgggtg	ctgatcgcca	cctccgtgga	gacatcagcc	gccagtggca	gccccgaggg	420
agctggaatg	accacagttc	agaccatcac	aggcagtgat	cccgagggaag	ccatctttga	480
caccttttgc	accgatgaca	gctctgaaga	ggcaaagaca	ctcacaatgg	acatattgac	540
attggtcac	acctccacag	aagctaaggg	cctgtcctca	gagagcagtg	cctcttccga	600
cggcccccat	ccagtcacac	ccccgtcacg	ggcctcagag	agcagcgcct	cttccgacgg	660
cccccatcca	gtcatcaccc	cgtcacgggc	ctcagagagc	agcgcctctt	cgcagggccc	720
ccatccagtc	atcaccccg	catgggtccc	gggatctgat	gtcactctcc	tcgttgaagc	780
cctggtgact	gtcacaaaac	tcgaggttat	taattgcagc	atcacagaaa	tagaaacaac	840
aacttccagc	atccctgggg	ctcagacat	agatctcact	cccacggaag	gggtgaaggc	900
ctcgtccacc	tcgatccac	cagctctgcc	tgactccact	gaagcaaaac	cacacatcac	960
tgaggtcaca	gcctctgccg	agacctgtc	cacagccggc	accacagagt	cagctgcacc	1020
tcatgccacg	gttgggaccc	cactccccac	taacagcgcc	acagaaagag	aagtgcacgc	1080
accggggg	acgacctca	gtggagctct	ggtcacagtt	agcaggaatc	ccctggaaga	1140
aacctagccc	ctctctgttg	agacaccaag	ttactgcaaa	gtctcaggag	cagctccggg	1200
ctccatagag	gctgggtcag	cagtgggcaa	aacaacttcc	tttgcaggga	gctctgcttc	1260
ctcctacagc	ccctcggaag	ccgcctcaa	gaacttcacc	ccttcagaga	caccgaccat	1320
ggacatcgca	accaaggggc	ccttccccac	cagcagggac	cctcttctct	ctgtccctcc	1380
gactacaacc	aacagcagcc	gagggacgaa	cagcacctta	gccaagatca	caacctcagc	1440
gaagaccacg	atgaagcccc	aacagccacg	cccacgactg	cccgagcag	gcccagcaca	1500
gacgtgagtg	caggtgaaaa	tggaggtttc	ctcctcctgc	ggctgagtg	ggcttccccg	1560
gaagacctca	ctgacccacg	agtggcagaa	aggctgatgc	agcagctcca	ccgggaactc	1620
cagccccacg	gcctcactt	ccaggtctcc	ttactgcgtg	tcaggagagg	ctaaccggaca	1680
tcagctgcag	ccaggcatgt	cccgtatgcc	aaaagagggg	gctgccccta	gcttgggccc	1740
ccaccgacag	actgcagctg	cgttactgtg	ctgagaggta	cccagaaggt	tcccatgaag	1800
ggcagcatgt	ccaagcccc	gacccagat	gtggcaacag	gaccctcgct	cacatccacc	1860
ggagtgtatg	tgtggggagg	ggcttcacct	gttcccagag	gtgtccttgg	actcaccttg	1920
gcacatgttc	tgtgtttcag	taaagagaga	cctgatcacc	catctgtgtg	cttccatcct	1980
gcattaaaat	tcactcagtg	tggcccaaaa	aaaaaa			2016

<210> 29
 <211> 2520
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1524355CB1

<400> 29						
caagatggcg	gctggcgagg	ctgtcgtctg	ggcgcccgag	tgcgggcttc	tcccctacgc	60
gctacacaag	tggagctcct	tttctctcac	ctaccttccc	gagaacattt	tagtggacaa	120
accaaagtac	caatcttcaa	gatggtcttc	agagagcaac	tatcctcccc	agtacttgat	180
tctaaagctc	gaaaggcctg	ctatagtcca	gaatatcaca	tttggaaaaa	atgagaaaaa	240
tcattgtttg	aatttgaaga	aatttaaagt	ctttgggtgga	atgaatgaag	aaaatatgac	300

WO 00/68380

PCT/US00/12811

agagctgttg	tccagtggct	ttaaagaatga	ttataacaaa	gaaacattca	ccttgaagca	360
taaaattgat	gaacagatgt	tccttctgtcg	attcattaaa	atagttccac	tcttgtcctg	420
gggacccagc	tttaacttta	gcattctggta	tggtgaaact	agtggcattg	atgacacctga	480
tatagtacaa	ccttgtctca	actggtatag	caagtaccgt	gaacaggaag	ctattcgcct	540
ttgcctaaaa	cacttcagac	aacacaaacta	tacagaagct	tttgagtcac	tgcaaaaagaa	600
aaccaagatt	gcactggaac	atcccatggt	aacagatatt	catgacaagc	tggtgttgaa	660
gggtgatttt	gatgcttgcg	aagagttgat	tgaaaaggct	gtaaatgatg	gcttgttcaa	720
tcagtatatc	agtcaacagg	aatataagcc	acgatggagt	caaatacttc	ccaaaagtaa	780
caaaggtgat	ggggaagata	accgtccagg	aatgagagga	ggccatcaga	tggttattga	840
tggtcaaaca	gagactgttt	atgtgtttgg	tggtctgggt	gggaacacaag	atcttctgtga	900
cttctgggcg	tacagtgtga	aggagaacca	gtggacatgt	atctctagag	acactgaaaa	960
agagaatggt	cctagtgtcca	gatcgtgtca	taaaatgtgc	attgatattc	aacggaggca	1020
aatctacaca	ttggggcggt	acttggtatc	ctctgtgagg	aacagcaaag	ctctgaaaag	1080
tgactttctat	cgttatgaca	ttgatacaaaa	cacatggatg	ttactaagtg	aggatactgc	1140
tgctgatgga	gggcccgaat	tggtgtttga	tcactcagatg	tgtatggact	cagaaaaaca	1200
tatgatctac	acttttgggtg	gtagaatttt	gacttgtaat	ggcagcgtag	ggcagcagag	1260
agccagtga	ccacaattca	gtggcttgtt	tgctttcaac	tgtcaatgtc	aaacctggaa	1320
acttcttcga	gaggactcct	gtaatgctgg	gcctgaggac	atccagtctc	gaataggaca	1380
ctgcatgtta	ttccactcaa	aaaatcgttg	cttatatgta	tttgggtggc	agcgatcaaa	1440
gacctattttg	aatgatttct	ttagttatga	tgtggactct	gatcatgtag	acataatatc	1500
agatggcacc	aagaaagact	ctgggatggg	tccaatgaca	ggattttacac	agagagcaac	1560
tattgatcca	gaactgaatg	aaatacacgt	cttatctgga	ctcagcaaag	ataaggaaaa	1620
gaggggaagaa	aatgttagaa	attcattctg	gattttatgac	attgtgagga	atagttgggtc	1680
ttgtgtctat	aagaatgata	aagctgcaaa	ggataatcca	actaaaagtc	ttcaggaaga	1740
agaaccatgt	ccaaggtttg	cccatcagct	tgtatacgat	gagctacaca	aggttcatta	1800
cttattttggt	gggaatccag	gaaaatcttg	ctctccaaag	atgagattag	atgactttctg	1860
gtcactgaag	ttgtgtagac	cttcaaaaaga	ttattttactg	aggcattgca	agtacctcat	1920
aagaaaacac	aggtttgaag	aaaaggccca	agtggtatccc	cttagtgctc	tgaaatattt	1980
acaaaatgat	ctttatataa	ctgtggatca	ttcagaccca	gaagagacaa	aagagtttca	2040
gctcctggca	tcagctctat	tcaaatctgg	ttcagatttt	acagctctgg	gctttttctga	2100
tgtggatcac	acctatgctc	aaagaactca	gctctttgac	accttagtaa	atctctttcc	2160
tgacagcatg	actcctccta	aaggcaacct	ggtagacctc	atcacactgt	aactgaagag	2220
tcactggaca	cagaaatgga	aaacaggagt	cgattttccg	tctttttggat	tgcagctcca	2280
ctgactgaca	gtaaagctgc	agtgattgag	gactgcacca	gagttctgaa	gggatcttaa	2340
ccatcacaaag	tttttaccct	cttccttcat	gcctgacctc	aaccccgctc	tcctcatcct	2400
attcctaata	taggctaata	aagtgaat	ggtatacttt	ccagttaaat	atatatatat	2460
atatattttt	tcttacttta	tcttttaaga	attaatgagt	ataaaagcaa	aattaggcag	2520

<210> 30

<211> 1954

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1598937CB1

<400> 30

ccgagttatg	cagctccggg	cggcaagggg	tcctgtcgag	gggcgcggtc	cataagggtt	60
gaggccagag	gcgcgtacct	ctggggcgccg	agctctggag	tggaggctgg	ggttcggagt	120
gcgtcatctg	gaagcaggca	ccccggcccg	caggcagagt	cacggtggca	gcattgagag	180
ttggacaccc	gggtccttga	agtgatctct	aggccccagc	cccaaatccg	ccaccattcc	240
gtgctgcggg	gacaccatgg	ctccagaaga	ggacgctgga	ggggaggcct	tagggggcag	300
tttctgggag	gctggcaact	acaggcgcac	ggtacagcgg	gtggaggacg	ggcaccggct	360
gtgcggggac	ctggctcagct	gcttccagga	gcgcgcgcgc	atcgagaagg	cttatgccca	420
gcagttggct	gactgggccc	gaaagtggag	ggggaccgtg	gagaagggcc	cccagtatgg	480
cacactggag	aaggcctggc	atgccttttt	cacggcggtc	gagcggctga	gcgcgctgca	540
ctgggaggtg	cgaggagaagc	tgcaagggca	ggacagttag	cgggtgcgcg	cctggcagcg	600
gggggctttc	caccggcctg	tgctggggcg	cttccgcgag	agccggggcg	ccgaggacgc	660
cttccgcaag	gcccagaagc	cctggctgaa	gaggctgaag	gaggttgagg	cttccaagaa	720
aagctaccac	gcagcccggga	aggatgagaa	gaccgcccag	acgagggaga	gccacgcaaa	780
ggcagacagc	gccgtctccc	aggagcagct	gcgcaaaactg	caggaacggg	tggaaacgctg	840
tgccaaggag	gccgagaaga	caaaagctca	gtatgacagc	acgctggcag	agctgcacgc	900
ctacactcca	cgctacatgg	aggacatgga	acaggccttt	gagacctgcc	aggcccgcca	960
gcgccagcgg	cttcttttct	tcaaggatat	gctgctcacc	ttacaccagc	acctggacct	1020
ttccagcagt	gagaagttcc	atgaactcca	ccgtgacttg	caccagggca	ttgaggcagc	1080
cagtgcagaa	gaggatctgc	gctgggtggcg	cagcaccac	gggccaggca	tggccatgaa	1140

WO 00/68380

PCT/US00/12811

ctggccacag	ttcgaggagt	ggtccttggg	cacacagagg	acaatcagcc	ggaaagagaa	1200
gggtggccgg	agccctgatg	aggttaccct	gaccagcatt	gtgcctacaa	gagatggcac	1260
cgcaacccca	ccccagtcct	cgggggtcccc	aggcacgggg	caggatgagg	agtggtcaga	1320
tgaagagagt	ccccggaagg	ctgccaccgg	ggttcgggtg	agggcactct	atgactacgc	1380
tggccaggaa	gctgatgagc	tgagcttccg	agcaggggag	gagctgctga	agatgagtga	1440
ggaggacgag	cagggctggt	gccaaggcca	gttgccagagt	ggccgcattg	gcctgtaccc	1500
tgccaactac	gtggagtgtg	tgggcgcctg	agtgtcctga	cagcccttct	gcaacgttta	1560
cccaccctgg	ttcagagccc	agcttctctc	ggagagccgg	acctcaggg	ccctgaaccg	1620
tcgctctctg	gctgctcctc	tgtcccttga	gggaggaagt	cctgggaccc	agggagggga	1680
ggggcctttg	tctagggag	ggactggtag	ggaagggacg	agtctaggct	gagggcaaga	1740
tgggaggtca	gaggtgacag	aagcgttcag	gggtgcctgg	gcctccccag	gagctgtgga	1800
ctcagttcct	gacctctgct	ttggggttcc	tggggtgggc	ttggggtgag	tgtagttctg	1860
gcctagcagc	acctctctgt	ggcttgttct	agcgtgtatt	aaaacttgac	acacacccac	1920
acacaaaaac	aaaaacacca	aaaaaaaaaa	aaaa			1954

<210> 31
 <211> 1817
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1725801CB1

<400> 31						
gacgcggtga	ggagacggcc	cacggcgccc	gcgggctggg	gcggtegett	cttccttctc	60
cggtggcctac	gaggggtcccc	agcctgggta	aagatggccc	catggccccc	gaagggccta	120
gtcccagctg	tgctctgggg	cctcagctc	ttcctcaacc	tcccaggacc	tatctggctc	180
cagccctctc	cacctcccca	gtcttctccc	ccgcctcagc	cccatccgtg	tcatacctgc	240
cggggactgg	ttgacagctt	taacaagggc	ctggagagaa	ccatccggga	caacttttga	300
ggtggaaaaca	ctgcctggga	ggaagagaat	ttgtccaaat	acaaagacag	tgagacccgc	360
ctggtagagg	tgctggaggg	tgtgtgcagc	aagtcagact	tcgagtgcc	ccgcctgctg	420
gagctgagtg	aggagctggg	ggagagctgg	tggtttcaca	agcagcagga	ggccccggac	480
ctcttcagtg	ggctgtgctc	agattccctg	aaagctctgt	gccccgcagg	ccacctcggg	540
ccctcctgcc	ttccctgtcc	tgggggaaca	gagaggccct	gcgggtggct	cgggcagtg	600
gaaggagaag	ggacacgagg	gggcagcggg	cactgtgact	gccaagccgg	ctacgggggt	660
gaggcctgtg	gccagtgtgg	ccttggctac	tttgaggcag	aacgcaacgc	cagccatctg	720
gtatgttcgg	cttgttttgg	cccctgtgcc	cgatgtctag	gacctgagga	atcaaactgt	780
ttgcaatgca	agaagggtct	ggccctgcat	cacctcaagt	gtgtagacat	tgatgagtgt	840
ggcacagagg	gagccaactg	tggagctgac	caattctgct	tgaacactga	gggctcctat	900
gagtgcagag	actgtgccaa	ggcctgccta	ggctgcattg	gggcagggcc	aggtcgctgt	960
aagaagtgtg	gccttggtta	tcagcaggtg	ggctccaagt	gtctcgatgt	ggatgagtgt	1020
gagacagagg	tgtgtccggg	agagaacaag	cagtgtgaaa	acaccgaggg	cggttatctc	1080
tgcatctgtg	ccgagggtta	caagcagatg	gaaggcatct	gtgtgaagga	gcagatccca	1140
gagtcagcag	gcttcttctc	agagatgaca	gaagacagag	tggtggtgct	gcagcagatg	1200
ttctttggca	tcatacatct	tgcactggcc	acgctggctg	ctaagggcga	cttggtgttc	1260
accgccatct	tcattggggc	tgtggcggcc	atgactggct	actggttgtc	agagcgcagt	1320
gaccgtgtgc	tggagggctt	catcaagggc	agataatcgc	ggccaccacc	tgtaggacct	1380
cctcccacc	acgctgcccc	cagagcttgg	gctgcctctc	tgctggacac	tcaggacagc	1440
ttggtttatt	tttgagagt	gggtaagcac	cctacactgc	cttacagagc	agccaaagta	1500
cccaggcccc	ggcagacaag	gcccctgggg	taaaaagtag	ccctgaaggt	ggataccatg	1560
agctcttcac	ctggcgggga	ctggcaggtc	tcacaatgtg	tgaatttcaa	aagtttttcc	1620
ttaatgggtg	ctgctagagc	tttggccctc	gcttaggatt	aggtggtcct	cacaggggtg	1680
gggccatcac	agctccctcc	tgccagctgc	atgctgccag	ttcctgttct	gtgttcacca	1740
catccccaca	ccccattgcc	acttatttat	tcattctcag	aaataaagaa	aggtcttgga	1800
aagttaaaaa	aaaaaaa					1817

<210> 32
 <211> 2694
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1730482CB1

<400> 32

gacctagtgt	gagcataatg	gaaaaaacac	aatcacttcc	tacacgacca	ccaacttttc	60
ctccaaccat	tccaccagca	aaagaagtat	gtaaggcggc	caaggtctgc	ctggtattta	120
tgggtggatg	atccctggagc	attggagatg	aaaatttcaa	taagatcatc	agcttttctat	180
acagcactgt	tggagccctg	aacaagattg	gcacagatgg	aacccaagtt	gcaatggttc	240
agttcactga	tgatcccaga	acagaattta	aactaaatgc	ttacaaaacc	aaagagactc	300
ttcttgatgc	aattaaacac	attttcatca	aaggaggaaa	tacaaaaaca	ggaaaagcaa	360
ttaatgatgt	tcgagatacc	ttgttctact	cagagtccgg	tacaagaagg	ggcatcccaa	420
aggttatcgt	gggtataact	gatggaagat	cacaagatga	tgtgaacaaa	atctccaggg	480
agatgcaatt	agatggctat	agcattttttg	caattggtgt	ggccgatgca	gattactcgg	540
agttgggttag	cattggcagt	aagcccagcg	cacgccatgt	cttcttttgt	gatgactttg	600
acgcctttaa	gaaaaatcgaa	gatgagctta	ttacttttgt	ctgcgaaaca	gcatcagcaa	660
ctgtccagt	ggtagacaaag	gatggcattg	atcttgcagg	atttaagatg	atggaaatct	720
ttgggttgggt	tgaaaaagat	ttttcatcag	tggaaaggggt	ttctatggag	ctgggtacct	780
tcaatgtgtt	tccatgtttac	caactccata	aagatgccct	ggttttccag	ccaaccaggt	840
acttgcaccc	agaaggattg	cctcccgact	acacaatcag	ttttctattc	cggattcttc	900
ctgacactcc	acaggagcca	tttgtctctt	gggagathtt	aaataaaaaa	ctgacccatc	960
tgggttgggggt	tatttttagac	aatggtggga	aaactctaac	atattttcaac	tatgaccaga	1020
gtgggggattt	tcaaactgttt	actttcgaag	gacctgaaat	taggaaaatt	ttttatggaa	1080
gctttcacaa	gctacacatt	gttgtcagtg	agacttttgtt	caaagtgggt	attgactgca	1140
agcaagtggg	tgagaaggca	atgaacgat	cagctaatat	cagctcagat	ggtgtagaag	1200
tgctagggaa	aatggttcga	tcaagaggac	caggtggaaa	ctctgcaccg	ttccagttac	1260
agatgtttga	tattgtttgc	tccacatcat	gggccaatat	agacaaatgc	tgtgaacttc	1320
caggcctgag	agatgatgag	tcttgcgccg	accttcccca	ttcctgtctc	tgtttctgaaa	1380
ccaatgaagt	ggctctggga	ccagcgggcc	caccaggtgg	tccaggactc	cgaggaccaa	1440
agggccagca	aggtgaaccg	ggtccaaagg	gaccagatgg	cctcgggggt	gaaattgggtc	1500
tgccaggacc	tcagggtcca	cctggacctc	aaggaccaag	tggctctgtcc	attcaaggaa	1560
tgcccggaat	gccaggagaa	aaaggagaga	aaggagatac	tggccttcca	ggtccacagg	1620
gtatcccagg	aggcgttggt	tcaccaggac	gtgatggctc	accaggccag	aggggccttc	1680
cgggaaagga	tggatcctcg	ggacctccag	gaccaccagg	gccaataggc	attcctggca	1740
ccctctggagt	cccagggatc	acaggaagca	tgggaccgca	aggcgccttg	ggaccacctg	1800
gtgtccctgg	agcaaaaggg	gaacgaggag	agcggggtga	cctgcagtct	caagccattg	1860
tgatcatcgt	ggcgcgtcaa	gtatgcgaac	agctcatcca	gagtcacatg	gccaggtaca	1920
ctgccatcct	caaccagatt	cccagccact	ctctatccat	cgggactgtc	caagggcctc	1980
ctggggagcc	tggggagcca	ggctcacctg	gagccccctg	tgaacaagga	cccccaggca	2040
caccaggctt	ccccggaaat	gcaggcgtgc	cagggacccc	agggagaacga	ggtcttaactg	2100
gtatcaaagg	agaaaaagga	aatccaggcg	ttggaaacca	aggtccaaga	ggccccctctg	2160
gaccagcagg	accttcaggg	gagagtggcg	ctggcagccc	tgggccccct	ggctctcctg	2220
gaccaagagg	cccaccagta	catctggggg	ttcctggacc	ccaaggctct	tctggccagc	2280
ctggtatattg	tgaccctcta	tcattgtctg	cctatgtgtg	gagagctccc	catccagatc	2340
agccagagtt	caccctctgc	caagatgagc	tggaaagccat	ggaactgtgg	ggccctggag	2400
tctgatagcc	tcaggagaaa	tttgaagacc	aactgcaaga	actcttaagg	aatcttgttt	2460
gagaaaaatct	tgttatgtgg	tttgatatgt	actttttggg	ggcagggtctc	atttcagcag	2520
cctaaatctc	ctccttggat	aatgttaata	ttattattat	tattaaacaa	aatatatatat	2580
tttataaaaag	ttcccttaat	ctatgacatg	gtagcaatga	tttccctttg	gtgtcttaat	2640
ggcatgtcag	ataattttgtt	tttccagaga	agagagctca	aagaggaatt	ggga	2694

```
<210> 33
<211> 1149
<212> DNA
<213> Homo sapiens
```

```
<220>  
<221> misc_feature  
<223> Incyte ID No: 1810058CB1
```

<400> 33						
cagtatcttg	gtccagcctg	cagccttagg	gtccagggtga	tgttacccgtg	tgtgtggccc	60
ttcttcacag	tggcctccta	gaaaaacaag	accctgactc	aaagaacacc	tctcactaca	120
ttcagagtct	gtcatactgaa	ccatgaggat	ctggtggcct	ctgcttgcca	tgaaatctg	180
cacagggaac	ataaactcac	aggacacctg	caggcaaggg	caccctggaa	tccctgggaa	240
ccccgtcac	aatggtctgc	ctggaagaga	tggacgagac	ggagcgaagg	gtgacaaagg	300
cgatgcagga	gaaccaggac	gtcctggcag	ccgggggaag	gatgggacga	gtggagagaa	360
gggagaacga	ggagcagatg	gaaaagt tga	agcaaaaggc	atcaaagggtg	atcaaggctc	420
aagaggatcc	ccaggaaaaa	atggccccca	ggggcttgca	gggcccattg	gagagaaagg	480
ctcccgagga	gagactgggc	ctcaggggca	gaagggggaat	aagggtgacg	tgggtccacc	540
tggctctgag	gggccaaagg	gcaacattgg	gcctttgggt	ccaacttggt	taccggggcc	600
catgggccct	attggaaagc	ctgggtccca	gggagaagct	ggaccacagg	ggccccaggg	660

WO 00/68380

PCT/US00/12811

atagttctag	ctctatcaga	ttcagtaaac	ttggatgaga	ttacattcca	catttgactc	960
tcagcttttag	agatatggta	acagaatttc	tacaacagat	cctgaattct	tattgcatta	1020
agggctctgc	tttggcttat	atgtgcatta	tcacacttaa	tcagtgcaa	cgtgccttta	1080
tcaccttgaa	gccagggtaa	acaaaggaag	agtgatttgc	atctaaagag	aacaaagccc	1140
caaccctctg	gctataccca	accactcaaa	ggcagcacag	gaacccacat	cactgcttgg	1200
ataatcccag	gaaaatgcag	aaaaagtgtg	gcctgaagca	tgattttctc	atgtggcact	1260
tctgtgtgca	ggagatcaca	gcgcggtttt	gttgctgcca			1300

<210> 36

<211> 1562

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3117318CB1

<400> 36

aaggccgggc	gcggttaagag	cgtctcgggg	agtagggcaa	ggcggccggg	cccccccat	60
tccgcctttt	cttcagcgtc	ctgcccggcg	cactggctgc	gggtgccggg	ccacctgcga	120
gtgtgctcag	ggactctgga	caccgcgggc	ggcgagctga	gggagcagtc	tccacgagga	180
cccaggcgga	ccctctggcg	ccatgcgcgc	cctccccggc	ctgctggagg	ccagggcgcg	240
tacgccccgc	ctgctcctcc	tccagtgcct	tctcgtctgc	gcgcgcccga	gctcggcgga	300
cggcagtgcc	ccagattcgg	cttttacaag	tcacactctc	agagaagaaa	taatggcaaa	360
taacttttcc	ttggagagtc	ataacatata	actgactgaa	cattctagta	tgccagtaga	420
aaaaaatatc	acttttagaaa	ggcctttctaa	tgtaaatctc	acatgccagt	tcacaacatc	480
tggggatttg	aatgcagtaa	atgtgacttg	gaaaaaagat	ggtgaacaac	ttgagaataa	540
ttatcttgct	agtgcacacag	gaagcacctt	gtatacccaa	tacaggttca	ccatcattaa	600
tagcaaacaa	atgggaagtt	attcttgttt	ctttcgagag	gaaaaggaac	aaagggggaa	660
atttaatttc	aaagtccctg	aacttcatgg	gaaaaacaag	ccattgatct	cttacgtagg	720
ggattctact	gtcttgacat	gtaaatgtca	aaattgtttt	cctttaaatt	ggacctggta	780
cagtagtaat	gggagtgtaa	aggttccctgt	tgggtgttcaa	atgaataaat	atgtgatcaa	840
tggaaacatat	gctaacgaaa	caaagctgaa	gataacacaa	cttttgagg	aagatgggga	900
atcttactgg	tgccgtgcac	tattccaatt	aggcgagagt	gaagaacaca	ttgagcttgt	960
ggtgctgagc	tatttggtgc	ccctcaaacc	atctcttgta	atagtggctg	aggtgattct	1020
tttagtggcc	accattctgc	tttgtgaaaa	gtacacacaa	aagaaaaaga	agcactcaga	1080
tgaggggaaa	gaatttgagc	agattgaaca	gctgaaatca	gatgatagca	atggtataga	1140
aaataatgtc	cccaggcata	gaaaaaatga	gtctctgggc	cagtgaatac	aaaacatcat	1200
gtcgagaatc	attggaagat	atacagagtt	cgtatttcag	ctttgtttat	ccttccctgtt	1260
aagagcctct	gagtttttag	ttttaaaagg	atgaaaagct	tatgcaacat	gctcagcagg	1320
agcttcacata	acgatatatg	tcagatctaa	aggtatatct	tcattctgta	attatgttac	1380
ataaaagcaa	tgtaaatcag	aataaatatg	ttagaccaga	atacaaatta	attatattct	1440
ggtcttcaaa	ggacacacag	aacagatata	agcagaatca	cttaataact	catagaacaa	1500
aatcactca	aaacctgttt	ataaccaaag	aattcatgaa	aaagaaagcc	tttggccatt	1560
tg						1562

<210> 37

<211> 2801

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> 2793

<223> a, t, c, g, or other

<220>

<221> misc_feature

<223> Incyte ID No: 3486992CB1

<400> 37

gacttttgctt	gaatgtttac	atcttctgct	cgtgttccta	catatcacaa	tatagtgttc	60
acgtttttgtt	aaaacttttg	ggtgtcagga	gttgagcttg	ctcagcaagc	cagcatggct	120
aggatgagct	ttgttatagc	agcttgccaa	ttgggtgctg	gcctactaat	gacttcatta	180
accgagtctt	ccatacagaa	tagtgagtgt	ccacaacttt	gcgtatgtga	aattcgctcc	240
tggttttacc	cacagtcaac	ttacagagaa	gccaccactg	ttgattgcaa	tgacctccgc	300
ttaacaagga	ttcccagtaa	cctctctagt	gacacacaag	tgcttctctt	acagagcaat	360

WO 00/68380

PCT/US00/12811

aacatcgcgga	agactgtgga	tgagctgcag	cagcttttca	acttgactga	actagatttc	420
tcccaaaaaca	acttttactaa	cattaaggag	gtcgggctgg	caaaccctaac	ccagctcaca	480
acgctgcatt	tggaggaaaaa	tcagattacc	gagatgactg	attactgtct	acaagacctc	540
agcaaccttc	aagaactcta	catcaaccac	aaccaaatta	gcactatttc	tgctcatgct	600
tttgcaggct	taaaaaatct	attaaggctc	cacctgaact	ccaacaaatt	gaaagtattt	660
gatagtcgct	ggttttgatc	tacacccaac	ctggaaattc	tcatgatcgg	agaaaaccct	720
gtgattggaa	ttctggatat	gaacttcaaa	ccccctgcaa	atttgagaag	cttagttttg	780
gcaggaaatg	atctcactga	tattccttga	aatgcttttg	tgggtcttga	tagccttgag	840
agcctgtctt	tttatgataa	caaactgggt	aaagtccctc	aacttgccct	gcaaaaagtt	900
ccaaattttga	aattcttaga	cctcaacaaa	aaccccatc	acaaaatcca	agaaggggac	960
ttcaaaaaata	tgcttcgggt	aaaagaactg	ggaatcaaca	atatgggcga	gctcgtttct	1020
gtcgaccgct	atgcccggga	taacttgcc	gaactcaca	agctggaagc	caccaataac	1080
cctaaactct	cttacatcca	ccgcttggt	ttccgaagt	tcctgtctct	ggaaagcttg	1140
atgctgaaca	acaatgcctt	gaatgccatt	taccaaaga	cagtcgaatc	cctccccaat	1200
ctgcgtgaga	tcagtatcca	tagcaatccc	ctcaggtgtg	actgtgtgat	ccactggatt	1260
aaactcaaca	aaaccaacat	ccgcttcctg	gagccccctg	ccatgttctg	tgccatgcg	1320
cccgaaatata	aagggcacca	ggtgaaggaa	gttttaatac	aggattcgag	tgaacagtgc	1380
ctcccaatga	tatctcacga	cagcttccca	aatcgtttaa	acgtggatat	cggcacgacg	1440
gttttccctag	actgtcgagc	catggctgag	ccagaacctg	aaatttactg	ggtcactccc	1500
attggaaata	agataactgt	ggaaacctgt	tcagataaat	acaagctaag	tagcgaaggt	1560
accttggaaa	tatctaacat	acaaattgaa	gactcaggaa	gatacacatg	tgttgcccag	1620
aatgtccaag	gggcagacac	tcgggtggca	acaattaagg	ttaacgggac	ccttctggat	1680
ggtacccagg	tgctaaaaat	atacgtcaag	cagacagaat	cccatcccat	cttagtgtcc	1740
tggaaagtta	attccaatgt	catgacgtca	aacttaaaat	ggctgtctgc	caccatgaag	1800
attgataacc	ctcacataac	atatactgcc	agggtcccag	tcgatgtcca	tgaatacaac	1860
ctaacgcac	tcgagccttc	cacagattat	gaagtgtgtc	tcacagtgtc	caatattcat	1920
cagcagactc	aaaagtcatg	cgtaaatgtc	acaacaaaaa	atgccgcctt	cgcagtggac	1980
atctctgatc	aagaaaccag	tacagccctt	gctgcagtaa	tggggctctat	gtttgcccgc	2040
attagccttg	cgtccattgc	tgtgtacttt	gcaaaaagat	ttaagagaaa	aaactaccac	2100
cactcattaa	aaaagtatat	gcaaaaaaac	tcttcaatcc	cactaaatga	gctgtaccca	2160
ccactcatta	acctctggga	aggtgacagc	gagaaagaca	aagatgggtc	tgcagacacc	2220
aagccaaccc	aggtcgacac	atccagaagc	tattacatgt	ggtaactcag	aggatatatt	2280
gcttctggta	gtaaggagca	caaagacgtt	tttgccttat	tctgcaaaag	tgaacaagtt	2340
gaagactttt	gtatttttga	ctttgctagt	ttgtggcaga	gtggagagga	cggttgagata	2400
tttcaaattt	ttttagtata	gcgtatcgca	agggtttgac	acggctgcca	gcgactctag	2460
gcttccagtc	tgtgttttgt	ttttattctt	atcattatta	tgattgttat	tatattatta	2520
ttttatttta	gttgttgtgc	taaactcaat	aatgctgttc	taactacagt	gctcaataaa	2580
atgattaatg	acaggatggg	gttccccctg	gcttttacca	gtagcatgac	cccttctgaa	2640
gccatccgta	gaaagtactt	tgtccccaaa	aagcaacata	cgggtttgaac	agcatgaaac	2700
tttgtagcat	cgggctaaga	ctttaactca	gagcaaggca	gactggtacc	tcgttaagat	2760
gtagtgactg	cggatgttta	cactgaatga	agntgcttaa	t		2801

<210> 38

<211> 2597

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4568384CB1

<400> 38

ccagaggcaa	agaggccagt	gaggactgct	ctgtgcagtt	gtccaggcct	gaagaagggtg	60
gcggtgattt	gaatcagaag	aggtggcatt	ctcttcatta	ggaggatatg	gataatcaag	120
acctaaacaaa	ttgggctccg	aaagatacct	tcgtcatggg	gaagagtga	tggagagaag	180
ataatataaaa	atgccaaacg	tatataaatt	gaaaggggaa	aattgagaga	acttgcgatc	240
catgggtcttg	gtcttccata	aaggggaact	gggacatcca	ctggagcaga	gcacagattg	300
gcccagagac	cccaagactc	ccactggcct	ccgcagaggc	cgacagtgt	ttcgctctgc	360
ggagattgtg	gcttccctgt	tagaaggaga	ggagaacacc	tgtggcaaac	agaaacaaa	420
ggaaaacata	ttaaagccaa	aatttcaggc	tttcaaggga	gtaggctgtc	tatatgaaaa	480
ggagtcaatg	aaaaaatcct	tgaaagacag	tgttgccctc	aacaataaag	atcagaattc	540
catgaaacat	gaggatccca	gtatcatatc	catggaagat	gggtccccat	atgttaattg	600
ctcattaggt	gaagtgactc	catgccaaac	tgcaaaaga	gcgaatggcc	caaactatat	660
tcagcctcaa	aaaagacaga	ccacttttga	aagccaggat	cgcaaggcag	gtccccctag	720
cagttctgaa	aagagaagta	agaatcctat	ttctaggcca	ttagaaggta	agaagtcctt	780
aagtcttagt	gcaaagactc	acaacatagg	ctttgacaaa	gacagctgcc	atagtaccac	840
aaagacagaa	gcttcacagg	aagagcgggc	tgattcaagc	ggcctcacat	ctctcaagaa	900

WO 00/68380

PCT/US00/12811

atcaccaaag	gtctcatcca	aggacactcg	ggaaatcaaa	actgatttct	cactttctat	960
tagtaattcg	tcagatgtga	gtgctaaaga	taagcatgct	gaagacaatg	agaagcggtt	1020
ggcagccttg	gaagcgaggc	aaaaagcaaa	agaagtgcag	aagaagctgg	tgcataatgc	1080
tctggcaaat	ttggatgggtc	atccagagga	taagccaacg	cacatcatct	tcggtttctga	1140
cagtgaatgt	gaaacagagg	agacatcgac	tcaggagcag	agccatccag	gagagggaatg	1200
ggtgaaagag	tctatgggtta	aaacatcagg	gaagctgttt	gatagcagtg	atgatgacga	1260
atctgattct	gaagatgaca	gtaatagggt	caaaattaaa	cctcagtttg	agggcagagc	1320
tggacagaag	ctcatggatt	tacagtcgca	ctttggcacc	gatgacagat	tccgcatgga	1380
ctctcgattt	ctagaaactg	acagtgaaga	ggaacaggaa	gaggtaaagt	aaaagaaaac	1440
tgctgaggaa	gaagagcttg	ctgaagaaaa	aaagaaagcc	ctgaatgttg	tacaaagtgt	1500
tttgcaaatc	aacttaagca	attctacaaa	cagaggatca	gtagctgcta	agaaatttaa	1560
ggacatcata	cattatgata	caacgaagca	agaccatgcc	acttacgaaa	gaaaaagaga	1620
tgataaacca	aaagaaagta	aagcaaaacg	aaaaaagaaa	agggaggaag	ctgagaaact	1680
acctgaggtg	tctaaagaaa	tgtattataa	tattgctatg	gatctgaaag	aaatatccca	1740
aactacaaaa	tataccagtg	aaaaggaaga	gggcacaccc	tggaaatgagg	actgtggtaa	1800
agacaaaact	gaggaaaacc	aggaccctgc	agctctgacc	agtgaagctg	agcagcccag	1860
cgggttcacg	ttctcttttt	ttgattcaga	cactaaagac	ataaagggaag	agacctacag	1920
agttgaaaca	gtgaaacctg	gaaagattgt	ctggcaggaa	gacctctggt	tacaagacag	1980
cagttcagaa	gaggaagatg	ttactgaaga	aacagatcac	agaaactcca	gtcctggaga	2040
agcatcatta	cttgagaaag	agaccactgt	atttttcttt	ttctctaaga	atgatgaacg	2100
acttcaaggt	tctgacttat	tctggagagg	agttaggaagt	aatatgagca	ggaactcttg	2160
ggaggccaga	acaaccaacc	tgcgtatgga	ttgtcgaaag	aaacataaag	acgcaaaaag	2220
gaaaatgaaa	ccaaaataat	aaatgtcagc	tggttttgat	actgaatgtg	aacaaggctc	2280
acctaaggaa	actgaccag	aaaacagttt	tagctgacaa	agaagaaatt	tcagagttaa	2340
ggaatttttaa	aaatctggct	gacggaatat	cattctgggt	gccatctttt	tctgtggaac	2400
tcctctgcat	ttcttcttaa	gtaattactt	caaaaattaa	attcaacttc	ttataaagga	2460
agaacaagat	agtccttgaa	aatacttttt	gtatataatc	tctttgcccc	ctatcctgag	2520
taactaatgg	acatcttctc	atgcaagggt	tatatgaagc	cttttttaaat	aaatgagtca	2580
aagcaaaaaa	aaaaaaa					2597

<210> 39

<211> 2641

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4586187CB1

<400> 39

ctgggaagaa	agctatcagc	accaactcag	aactctccac	cttcagatca	gacattctag	60
atctccgtca	gcaacttcgt	gagattacag	aaaaaaccag	caagaacaag	gatacgctgg	120
agaagttaca	ggcgagcggg	gatgctcttg	tggacaggca	gagtcaattg	aaagaaactt	180
tggagaataa	ctctttcctc	atcaccactg	taaacaaaac	cctccaggcg	tataatggct	240
atgtcacgaa	tctgcagcaa	gataccagcg	tgctccaggg	caatctgcag	aaccaaagt	300
attctcataa	tgtggtcatc	atgaacctca	acaacctgaa	cctgaccag	gtgcagcaga	360
ggaacctcat	cacgaatctg	cagcggctctg	tggatgacac	aagccagggt	atccagcgaa	420
tcaagaacga	ctttcaaaa	ctgcagcagg	tttttcttca	agccaagaag	gacacggatt	480
ggctgaagga	gaaagtgcag	agcttgccag	cgctggctgc	caacaactct	gcgttggcca	540
aagccaacaa	cgacaccctg	gaggatatga	acagccagct	caactcattc	acaggtcaga	600
tggagaacat	caccactatc	tctcaagcca	acgagcagaa	cctgaaagac	ctgaggact	660
tacacaaaga	tgcagagaat	agaacagcca	tcaagttcaa	ccaactggag	gaacgcttcc	720
agctctttga	gacggatatt	gtgaacatca	ttagcaatat	cagttacaca	gcccaccacc	780
tgcggagcgt	gaccagcaat	ctaaatgaag	tcaggaccac	ttgcacagat	acccttacca	840
aacacacaga	tgatctgacc	tccttgaata	ataccctggc	caacatccgt	ttggattctg	900
tttctctcag	gatgcaacaa	gatttgatga	ggtcagaggt	agacactgaa	gtagccaact	960
tatcagtgat	tatggaagaa	atgaagctag	tagactccaa	gcatggctag	ctcatcaaga	1020
atttttacaa	actacaaggt	ccaccgggccc	ccagggggtcc	aagagggtgac	agaggatccc	1080
agggaccccc	tggcccaact	ggcaacaagg	gacagaaagg	agagaagggg	gagcctggac	1140
cacctggccc	tgcgggtgag	agaggcccaa	ttggaccagc	tggtccccc	ggagagcggtg	1200
gcggcaaagg	atctaaaggc	tcccaggggcc	ccaaaggctc	ccgtgggttcc	cctgggaagc	1260
ccggccctca	gggccccag	ggggaccag	gccccccggg	cccaccaggc	aaagagggag	1320
ccccggccc	tcagggccct	cctggcttcc	agggacttca	gggcaccgtt	ggggagcctg	1380
gggtgctgg	acctcgggga	ctgccaggct	tgcttgggt	accaggcatg	ccaggcccca	1440
agggcccccc	cgccctcct	ggcccatcag	gagcgggtgt	gcccctggcc	ctgcagaatg	1500
agccaacccc	ggcacccggg	gacaatagct	gcccgcctca	ctggaagaac	ttcacagaca	1560
aatgctacta	tttttcagtt	gagaaagaaa	tttttgagga	tgcaaaagctt	ttctgtgaag	1620

WO 00/68380

PCT/US00/12811

tcacggcaac	agcatccagc	gcctggggga	ggtgaataag	ctggctgtcc	ttcctcggct	720
cctagcctg	acatccatg	ggaacccat	ggaggaagag	aaagggtata	ggcaatatgt	780
gctgtgcacc	ctgtcccgta	tcaccacggt	cgacttcagt	ggggtcacca	aagcagaccg	840
caccacagct	gaagtctgga	aacgcataaa	catcaagccc	aagaaggcct	ggaccaagca	900
gaatacactt	tgaggctccc	acgaccctag	tagtcctaaa	ggcctaagca	tagacagcat	960
ggtttgacaa	taaataattt	gagctgttga	gcagaaaaaa	aaaaaa		1006

<210> 42
 <211> 2582
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1833221CB1

<400> 42

gttaaatTTA	gtactgaatg	cagccttttg	ttgtttaaaa	aatttttttg	aactcacagt	60
ctacatcagc	atcagcatct	gcgtcaccat	ttcaatctgc	atgggtatagt	gaatctgaga	120
taactcaggg	agcacgctca	agatcgacga	accagcaacg	ggatcatgat	tcaaaaagac	180
ctaaactttc	ctgtacaaac	tgtactacct	cagctgggag	aatgtttgga	aattggttaa	240
acacattatc	agattcatct	tggaggcata	gtcaagttcc	tagatcttca	tcaatggtac	300
ttggatcatt	tggacacagc	ttaatgagag	agaggagaga	tttggagaga	agaacagatt	360
cctctattag	taatcttatg	gattatagtc	accgaagtgg	tgatttcaca	acttcatcat	420
atgttcaaga	cagagttcct	tcatattcac	aaggagcaag	accaaaagaa	aactcaatga	480
gcactttaca	gttgaataca	tcatccacaa	accaccaatt	gccttctgaa	catcagacca	540
tactaagttc	tagggactcc	agaaattcct	taagatcaaa	tttttcttca	agagaatcag	600
aatcttcccg	aagcaatacg	cagcctggat	tttcttacag	ttcaagtaga	gatgaagccc	660
caatcataag	caattcagaa	aggggtgttt	catctcaaa	accatttcaa	gaattctctg	720
acaatgaagg	taggcggaca	acgaggagat	tgctgtcacg	catagcttct	agcatgtcat	780
ctactttttt	ttcacgaaga	tctagtcagg	attccttgaa	tacaagatca	ttgaattctg	840
aaaattctta	cgtttctcca	agaatcttga	cagcttcaca	gtcccgtagt	aatgtacctt	900
cagcttctga	agttcccgat	aatagggcat	ctgaagcttc	tcagggattt	cgatttctta	960
ggcgaagatg	gggtttgtca	tctcttagcc	acaatcatag	ctctgagtca	gattcagaaa	1020
atTTtaacca	agaatctgaa	ggtagaaata	caggaccatg	gttatcttcc	tcacttagaa	1080
atagatgcac	acctttgttc	tctagaagga	ggcgagaggg	aagagatgaa	tcttcaagga	1140
tacctacctc	tgatacatca	tctagatctc	atatTTtttag	aagagaatca	aatgaagtgg	1200
ttcaccttga	agcacagaat	gatacctctg	gagctgtctg	caacagacca	caagcatctg	1260
cagcatcaag	cagtgccaca	acaggtggct	ctacatcaga	ttcgggtcaa	ggtggaagaa	1320
atacaggaat	atcagggatt	cttctgtggt	ccttattccg	gtttgcagtc	cccccagcac	1380
ttgggagtaa	tttgaccgac	aatgtcatga	tcacagtaga	tattattcct	tcagggttga	1440
attcagctga	tggtaaaagt	gataaaaact	aaagtgcgcc	ttcaagagat	ccagaaagat	1500
tgcagaaaat	aaaagagagc	ctccttttag	aggactcaga	agaagaagaa	ggtgacttat	1560
gtagaatttg	tcaaatggca	gctgcacatc	catctaattt	gctgatagag	ccatgcaagt	1620
gcacaggaag	tttgagtagt	gtccaccaag	actgtatgaa	aaagtgggta	caggccaaaa	1680
ttactcttgg	ttcttcatta	gaagctgtaa	ccacctgtga	actatgtaaa	gagaagttgg	1740
agcttaacct	ggaggatttt	gatattcatg	aactacatag	agctcatgca	aatgaacaag	1800
ctgagtatga	gtttatcagc	tctgggtctc	acctagtggg	gttattgcac	ttgtgcgaac	1860
aaagcttttc	tgatatgatg	ggaaatacaa	atgaaccaag	cacacgtgtc	cgatttatta	1920
accttgcaag	aactcttcag	gcacatatgg	aagatctcga	aacttcagag	gatgattccg	1980
aagaagacgg	agaccataac	aggacatttg	atattgccta	acttcatata	agacagatgg	2040
atgatctgtg	aacataaagt	tttattaaaa	atggcaatta	aatataaatt	acttttgtgg	2100
gggaatgcct	aataaataca	ttgactatat	ataaaatgaa	tatatacata	cacatgtatg	2160
cctgtatata	tatattcatt	ctccagtgtt	gctgaattaa	aattctgtctg	gactttttta	2220
catagcaaat	ccgatgttta	taaactggta	atcaaaaagg	ttttttcttt	taggtgagtg	2280
ggaaagtatt	acccttggtt	taaatatcta	agcaatgcct	atcaaccctt	ttttgtgtta	2340
tgattactgt	agtcataatt	atgaaaaaag	gtttgtgttt	tactcttgct	agtgagaaaa	2400
gtgggacaaa	atatactttt	gaaataaaaa	gctatatggc	acctaattat	tttttctttt	2460
aaaatgcctt	aagtgtcagt	ctcattttga	taatcatttg	cttcagtggt	ttaaaaatta	2520
aaaaaagaat	ggggagaagg	ttatgagaag	agcattatta	agtttccaaa	tttaatttga	2580
at						2582

<210> 43
 <211> 2849
 <212> DNA
 <213> Homo sapiens

WO 00/68380

PCT/US00/12811

<220>
<221> misc_feature
<223> Incyte ID No: 2041168CB1

<400> 43

```
acgtgaccaa gagctgacgt gtgcagaagt ccttcttctg ctggctcgttg ttcccgtctg 60
agtagcagct cccactgcc ctgagggcgg gccggcctgc ggcggaggga aaaaggaaga 120
ggagaaggaa attgtcccgga atccctgcag tctttctgta ggttgccggca caacgccagg 180
caaaagaaga ggaaggaatt taatcctaata cgggtggaggt cgatttgagg gtctgctgta 240
gcaggtgggt ccgcttgaag cgagggagga agtttccctcc gatcagtaga gattggaaag 300
attgttggga gtggcacacc actagggaaa agaagaaggg gcgaactgct tgtcttgagg 360
aggccaaccc ccagaatcag ctcttctggc cttgaagtgg ctgaagacga taccctcca 420
caggcttgag cccagtecca cagccttccct cccccagcct gagtgaactac tctattcctt 480
ggctccctgct attgtcgggg acgattgcat gggctacgcc aggaaagtag gctgggtgac 540
cgcagcctgg tgattggggc tggcgccctgc tattgcattt atagactgac taggggaaga 600
aaacagaaca aggaaaaaat ggctgagggg ggatctgggg atgtggatga tgctggggac 660
tgttctgggg ccaggtataa tgactgggtct gatgatgatg atgacagcaa tgagagcaag 720
agtatatgat ggtacccacc ttgggctcgg attgggactg aagctggaac cagagctagg 780
gccagggcaa gggccagggc taccggggca cgtcgggctg tccagaaacg ggcttcccc 840
aattcagatg ataccgtttt gtccctcaca gactataaaa aggttctttg cttgggttag 900
atgtctgaaa agccttatat tcttgaagca gctttaattg ctctgggtaa caatgctgct 960
tatgcattta acagagatat tattcgtgat ctgggtgggt tcccaattgt cgcaaagatt 1020
ctcaatactc gggatcccat agttaaggaa aaggctttaa ttgtcctgaa taacttgagt 1080
gtgaatgctg aaaatcagcg caggcttaaa gtatacatga atcaagtgtg tgatgacaca 1140
atcactttct ccttgaactc atctgtgcag ttgtctggac tgagattgct tacaaatatg 1200
actgttacta atgagtatca gcacatgctt gctaattcca tttctgactt tttctgttta 1260
ttttcagcgg gaaatgaaga aaccaaactt caggttctga aactcctttt gaatttggct 1320
gaaaatccag ccatgactag ggaactgctc agggcccaag taccatcttc actgggctcc 1380
ctctttaata agaaggagaa caaagaagtt attcttaaac ttctgggtcat atttgagaac 1440
ataaatgata atttcaaatg ggaagaaaat gaacctactc agaatcaatt cgggtgaagg 1500
tcactttttt tcttttttaa agaatttcaa gtgtgtgctg ataaggttct gggaatagaa 1560
agtcaccatg attttttggt gaaagtaaaa gttggaaaat tcatggccaa acttgctgaa 1620
catatgttcc caaagagcca ggaataacac cttgattttg taatttagaa gcaacacaca 1680
ttgtaaacta ttcattttct ccaccttggt tatatggtaa aggaatcctt tcagctgcca 1740
gttttgaaata atgaatatca tattgtatca tcaatgctga tatttaactg agttgggtctt 1800
taggttttaag atggataaat gaatatcact acttgttctg aaaacatggt tgttgctttt 1860
tatctcgctg cctagattga aatattttgc tttcttctct gcataagtga cagtgaacca 1920
attcatcatg agtaagctcc cttctgtcat tttcattgat ttaattttgt tatcatcaat 1980
aaaattgtat gttaatgctg gaaagaaaaa aagaagaaag aaaaaaacca tccctgtcct 2040
tcagtttata atctagttgg agagataaga aacgtacaaa ccaaaagata acagaatatc 2100
tgaagctagt actcattgtc agatgttccc tctgagagca cagaggaggc aaaagcttct 2160
gtgggatgtg ctagtcggct aaagcttcac agaggagggt gcaattgaaa atgagtcctg 2220
aatggggtag ggtggttagg gaattccatg agacaagaca aggggggcat ggtgtgagaa 2280
aggcatggaa gtaggaaccc tcttccatag acaggagatc attctgctta gagtggagag 2340
tgtggagagt gggagtagat aatttttgaa agctgggtga agccagttgt ggagaattgt 2400
ttgaattatta tcccattgaa taccagagac cactaaaatc ttttttacta gaaaataatt 2460
ggggtccata tgaaagtctc tattactgag tagtgtcaat gaggggtgtg caaaatgatt 2520
cctttcacat cctagtgggt gccatttggt aatacagata taagccttaa actatgtaaa 2580
cccttgctct aaggaagtaa ttgaataatt gcccaaagat tgtatgtatg aggtgttca 2640
tcccagcact gtctaagcta gtaaaaattg gaaacaattt aagtatctag cacattggat 2700
tggttataaa gcaaggaatg ttcacacagt aggatattat aagtatgctg atggaaatct 2760
atattgccag gaaaagctat tcattatgct ttgtgaagtc agaaagtaaa aaagggtaga 2820
tagaagtatt cgaagtatag ttccatttt 2849
```

<210> 44
<211> 670
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2365794CB1

<400> 44

```
gaggcaagaa ttcggcacga gggcgccgc gggcatttct tccactgccc gtctgaggga 60
acgctaagta gtgtgtccgg cgccgtgttc cagctccgcg ttgttccgcg agaaagcgag 120
aggccgagcc cgggctggtg cgatggccgc ggtggtggcc aagcgggaag ggccgccgtt 180
```


WO 00/68380

PCT/US00/12811

<221> misc_feature

<223> Incyte ID No: 2622288CB1

<400> 46

```
gcagaggcgg cggggctect cctcccgcgc ctctctcgcc tccccttcgg gcgctctcgc 60
gctaactgtg ctctctccgg gccctccgcc tgctcccage catggtggcc tggcgctcgg 120
cgttccttgt ctgcctcgct ttctccttgg ccacctggg ccagcgagga tctggggact 180
ttgatgattt taacctggag gatgcagtga aagaaacttc ctcatgaaag cagccatggg 240
accacaccac caccaccaca accaataggc caggaaaccac cagagctccg gcaaaacctc 300
caggtagtgg attggacttg gctgatgctt tggatgatca agatgatggc cgcaggaaac 360
cgggtatagg aggaagagag agatggaacc atgtaaccac cagcaccag aggccagtaa 420
ccaccagagc tccagcaaat actttaggaa atgattttga cttggctgat gccctggatg 480
atcgaaatga tcgagatgat ggccgcagga aaccaattgc tggaggagga ggtttttcag 540
acaaggatct tgaagacata gtaggggggt gagaatacaa acctgacaag ggtaaagggtg 600
atggccgggt aggcagcaat gacgacctcg gatctggcat ggtggcagag cctggcacca 660
ttggccgggt ggccagcgcc ctggccatgg cctcatcggg tgccgtctcc agctacatct 720
cctaccagca gaagaagttc tgcttcagca ttcagcaggg tctcaacgca gactacgtga 780
agggagagaa cctggaagcc gtggtatgtg aggaacccca agtgaataac tccacgttgc 840
acacgcagtc tgcagagccg ccgcccgcgc ccgaaccage ccgatctga gggccctgtc 900
cagctgcagg catgcacaat ggtgccaccg ctgtgccacc ggctccccc acccttcat 960
ttggacccgc agctgctgtg ctgctctgtg ccacgcgctc cttgttgggtc tgagtttccc 1020
ggatgagctc tgggtgtttg tgagtttggg ttctctgccc tgccccaagc gtgctgagac 1080
ttgggtgcga aattcaagag ccagctctga tagaaagcca gcaccagcct cgggagctgc 1140
tgagccacca actcccaaag ccagcctgcc tccagcttta ctgagcacag gatcggggg 1200
ccaagatgat gctgaggcct gatgacattt atgcttaggg gacaagagtt tgaactcaag 1260
ggactgtgac cctgcacac tggagtggct cattgtggca ggtttctgcc aatagacagc 1320
cctgcagagt ggctcaagg agctgcaggt ggggggctca gcctgcaccc acttggagcc 1380
cctgcaagga tgcgaacggg cagcaccag gttttatggt acaacatgc atctcgtgat 1440
atggtttcac ttcagtcttc cccatcccag gttttatggt acaacatgc atctcgtgat 1500
gtccaagcgg aggttttcag tgattttaagt ggataagtat gaagcaattc gtaggcctgt 1560
tgagagctta ggaatcttcc gggtctgggt tggccttcgg ttgacatga gaaaggtctt tagcaatcat 1620
attccatagg gggtctgggt gagatgagtt ttatcctgtg ttggggagag gtgctcacc tccacctgt 1680
ttctgcaccg ggttagcaag agtgaccgat gtcaagaacg agcatcaaag ccagaatcct 1740
gtccctgttt ttaaaaatgt aattgggggc ggcgggggag gagaggggaa agagacattc 1800
gcttgtttgc gtgaaacgca ggtgactttg tagctctgtg gtcagcctac ttgtctgctc 1860
gcttgggtta tgogtgggga gccatgctca ccgtggcaca cacaggaacc ccatgactcg 1920
gcttgggtta tgogtgggga gccatgctca ccgtggcaca cacaggaacc ccatgactcg 1980
cccctcacct ggctgggagc tgctgtgttt gggctggagc agagctgggt tcttggaatg 2040
ttcctttggc ccacatatgg ttctgtcccg gtgagctctg ttgtcagagg ctacacgggac 2100
agaaccacat gctaggtgtc agggcccttg tctactgata gtcagtttgc tgtgtcagaa 2160
agcacttctg aaagcagata tgagtcacca gacaggcagg atcttacaaa actcagggg 2220
ctctttgggtc tgcatgatgg ccccatgcgt ttcataggc gtcactgag cgggatgtgc 2280
tgctgagtg gatgagccaa ctccagtttc ttaaggaaac cactggaatc tgcagcccc 2340
acatgcactc gtctaacgca tgctctgtg tctgttttgc aacatgcctg tgggtggagg 2400
tgatgacttg tagcctgtg cgtctcaagg ctgcctgtg aggccattcc cagtgcgtgc 2460
ccttgagctc cttaccaccc cttttcctgc tccgcccttt aatccctgac agacctggac 2520
tgtgtggctg aagggggacc tgcagcactg cagaaatgcc tctgcgtggt gccatgaagg 2580
aaagaaacct tggcctgggt tgcagaagct tcccatgctt caggaaagtt gtaagggtgg 2640
ggtggctagg aggatggcc cctgatacca gctctccaca ctcatggcc agattgtgaa 2700
ctttgtcagg cttgtccctc cctgatacca gctctccaca ctcatggcc agattgtgaa 2760
tggctgggtc tgggcccggg gtggtatggg tttaccttta tgggtaggcc tgtcggatgg 2820
ctggacagcg tgacctcagg ggttgtccac ttagaatatg gctgtctgtg agacctgtt 2880
ctaagtcctt gaaaccctag agctgtgacg cccattttgc tctcccttac cgcccccg 3000
cccaggagca ctgactgcag ttgagagaga ctggcactca agcccccg tggcagctc 3060
cccggtgtc acatgcctgg aagaccctc agcctctgt ttgcagaacg ttcatttcag 3120
tagtgactgg ttcccacaga catcttacac ttgctcgaca ctgccacctg cagaagcctg 3180
gagcttctcc gtcacctgt gtcacttga aggttgact ggccagcatg ggcctgtccc 3240
aagcgagagg ggagacacag tggactgaaa ggaactgggt aaagtggcca atctctatca 3300
gcttaatttg gcagagaaaa tttgtaacaa ctctgagcac atgctgggtg aagtcacagc 3360
tcaaggaaag ataaagctg gcggaaggag gtgtgcgtgg cttctggggg gggaccaga 3420
ggggaggctc tgggacaggg gctgggggtt agtgccaggg cctgaggaa gaaatgggga 3480
ctgatctcaa aattccagaa ttccctgtac atctgttgc gtgcttgtg caaggtgtga 3540
cttgtaaaact gtctagtgtt tgcattaaat aaaatggcac cgagcagaaa aaaaaaaaa 3600
```

<210> 47

<211> 1236

WO 00/68380

PCT/US00/12811

<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2806595CB1

<400> 47
ttaaattccc cgaaatcaga ctgctgcctt ggaccgggac agctcgcggc ccccgagagc 60
tctagccgtc gaggagctgc ctggggacgt ttgccctggg gcccagcct gcccgggtc 120
accctggcat gaggagatgg gectgttget cctgggtccc tgctcctgc tgccgggtc 180
ctacggactg ccttcttaca accgttctta ctactccaac agcgccaacg accagaacct 240
aggcaacggt catggcaaaag acctccttaa tggagtgaag ctgggtgggtg agacacccga 300
ggagaccctg ttcacctacc aagggggccag tgtgatcctg ccctgccgt accgctacga 360
gccggccctg gtctccccgc ggcgtgtgcg tgtcaaatgg tggaaagctgt cggagaacgg 420
ggccccagag aaggacgtgc tgggtggccat cgggctgagg caccgtcct ttggggacta 480
ccaaggccgc gtgcacctgc ggcaggacaa agagcatgac gtctcgtggt agatccagga 540
tctgcggctg gaggactatg ggcgttaccg ctgtgaggtc attgacgggc tggaggatga 600
aagcgttctg gtggagctgg agctgcgggg tgagatgcta acggggactg ggtgacactg 660
ggacctgaga gcagagggga gaggaccaga gaaaacatcc agacctctgt gctttagaca 720
tttaaaaggt ctttaattctc aaaacaaccc acatgggaag tactgttgtg accccattt 780
tacaggtgag aaaactgagg cacagagagg tcaagtaact tacctaaggt cacacagctt 840
gtaaacgaca gagctgggggt ctgaacccaa gcaccagcc tctagaatct gtccccctct 900
accgctgtga attcacatct cattcagaga gaggaaaacc agagctggct ccacagctta 960
ttagagacag agctgagatt taagcaaggt tagcgggtaa cacaagcgaa tgaggcagcc 1020
cactgtgaga cgatgggagt ggtggggact gggcacactc ctgagccctt gtctgtgcc 1080
cagggagctc ccacactact caggcccagc tgatgatgcc acgccagaat gcagacccca 1140
ctgccaaatt ttctggcttg acaagataag cttatatatt tatgtgagat ctccagattt 1200
ttatgtaaaa acctcctttt taaaaacaaa acaaaa 1236

<210> 48
<211> 3081
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2850987CB1

<400> 48
gcccggccca cggcgggcgc ggcggggcgc gagagagctg gctcagggcg tccgctaggc 60
tcggacgacc tgcctgagct cccaaaccgc ttccataagg ctttgccttt ccaacttcag 120
ctacagtgtt agctaagttt ggaaagaagg aaaaaagaaa atccctgggc ccttttctt 180
ttgttctttg ccaaagtcgt cgttgtagtc tttttgccc aggtgttgt gtttttagag 240
gtgctatctc cagttccttg cactcctgtt aacaagcacc tcagcgagag cagcagcagc 300
gatagcagcc gcagaagagc cagcgggggc gctagtgtc atgaccaggg cgggagatca 360
caaccgccag agaggatgct gtggatcctt ggccgactac ctgacctctg caaaattcct 420
tctctacctt ggtcattctc tctctacttg gggagatcgg atgtggcact ttgcgggtgtc 480
tgtgtttctg gtagagctct atggaaacag cctccttttg acagcagctc acgggctggt 540
ggtggcaggg tctgtttctg tctgtggagc catcatcggt gactgggtgg acaagaatgc 600
tagacttaaa gtggcccaga cctcgttgtt ggtacagaat gtttcagtca tctgtgtgg 660
aatcatcctg atgatggttt tcttacataa acatgagctt ctgacctgt accatggatg 720
ggttctcact tctgtctata tctgatcat cactattgca aatattgcaa atttggccag 780
tactgtact gcaatcacia tccaaaggga ttggattgtt gttgttgcag gagaagacag 840
aagcaaacta gcaaatatga atgccacaat acgaaggatt gaccagttaa ccaacatctt 900
agccccatg gctgttggcc agattatgac atttggctcc ccagtcacg gctgtggctt 960
tatttcggga tggaaacttg tatccatgtg cgtggagtac gtctgtctc ggaagggtta 1020
ccagaaaacc ccagctctag ctgtgaaagc ttgtcttaaa gaagaggaaa ctgaattgaa 1080
acagctgaat ttacacaaag atactgagcc aaaaccctcg gagggaaact atctaattggg 1140
tgtgaaagac tctaacatcc atgagcttga acatgagcaa gagcctactt gtgcctccca 1200
gatggctgag ccttccgta ccttccgaga ttgatgggtc tctactaca accagcctgt 1260
gtttctggct gtcattgggt ttgctttcct ttatatgact gtctgtgggt gtcactgtat 1320
caccacaggg tacgcttaca ctacgggact gagtgggtcc atctcagta ttttgatggg 1380
agcatcagct ataactggaa taatgggaac tgtagctttt acttggctac gtcgaaaatg 1440
tggtttggtt cggacaggte tgatctcagc attggcacag ctttctgtgt tgatcttgtg 1500
tgtgatctct gtattcatgc ctggaagccc cctggaacttg tccgtttctc cttttgaaga 1560
tatccgatca aggttcattc aaggagagtc aattacacct accaagatac ctgaaattac 1620

WO 00/68380

PCT/US00/12811

<210> 50
<211> 1712
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 4675668CB1

```

<400> 50
ctttcttcag tccccacgtg cgatccttcc cggcaacttt ttcgagaaaa atgcccaaat 60
tcaaggcggc ccgtggggtg gggggtcagg aaaaacatgc gccctgggcc gatcagatcc 120
tggctgggaa tgcggtgcgg gcgggggtcc gggagaagcg gcggggtcgc gggacaggag 180
aagcggagga agagtatgtg gggccccggc tgagccgacg gattttgcag caagcacggc 240
agcaacagga ggaactcgag gccgagcatg ggactgggga caagcccgcg gcgccgcggg 300
aacgcaccac gcggctgggt ccaagaatgc ctcaggatgg atcagatgac gaggacgagg 360
agtggcccac cctggagaag gctgccacaa tgacagcagc gggccatcat gcagaggtgg 420
ttgtggaccc tgaggatgag cgtgccatag agatgttcat gaacaagaac cctcctgcca 480
ggcgacacct ggctgacatc atcatggaga agctgactga gaagcagaca gaggttgaga 540
cagtcatgtc agaggtgtcg ggcttcccta tgcccagct ggaccccccg gtcctagaag 600
tgtacagggg ggtccgggag gtattatcta agtaccgcag tggaaaactg cccaaggcat 660
ttaagatcat ccctgcactc tccaactggg agcaaatect ctacgtcaca gagccggagg 720
cctggactgc agctgccatg taccaggcca ccaggathtt tgccctaac ctgaagggaac 780
gcatggccca gcgcttctac aaccttgctc tgctccctcg agtacgagat gacgttgctg 840
aatacaaacg actcaacttc catctctaca tggctctcaa gaaggccctt ttcaaacctg 900
gagcctgggt caaagggatc ctgattccac tgtgcgagtc tggcacttgt accctccggg 960
aagccatcat tgtgggtagc atcatcacca agtgtcccat ccctgtgttg cactccagtg 1020
cggccatgct gaaaattgct gagatggaat acageggtgc caacagcatc ttcttgcgac 1080
tgctgctgga taagaagtat gcactgcctt accgggtgct ggatgcccta gtcttccact 1140
tcctgggggt ccggacagag aagcgtgaac tgccctgtgt gtggcaccag tgcctcctga 1200
ctttggtcca gcgtacaag gccgacttgg ccacagacca gaaagaggcc ctcttagaac 1260
tgctcgggct gcagcccat ccacagctat cgcccgaaat caggcgtgag cttcagagtg 1320
cagtcccccg cgatgtggaa gatgttccca tcaccgtgga gtgaggaaaa cagtcagctg 1380
tcctggccaa aggggttttg aaggacacca agacccccgt tgggtgactga agatgacact 1440
gagctttaat ggctgaagac ccagatcagg gcagtgacag atcacaggga catctgtggc 1500
tcccagtcca ggacaggaag gactgagggg ctggctgggt ccctcttcca ttctaggccc 1560
ttatccctgt ttagttctga gagccaactt gagataccat atgctagcat tccagtcctc 1620
cagctggggc ttggtgtgag tactttttct atggctattg tgtcagggtc ctgtggataa 1680
aggcaaaagc agatatttat tgaaaaaaaa aa 1712

```